

The Materia Medica of the Nosodes.

By Henry Clay ALLEN, M. D.

Materia Medica of the Nosodes (Only Key-Notes).

Ambra grisea * Anthracinum * Electricity * Lac caninum * Lac vaccinum * Lac vaccinum defloratum *
Lyssin * Magnetis poli ambo * Magnetis polus arcticus * Magnetis polus australis * Malandrinum *
Medorrhinum * Psorinum * Secale cornutum * Syphilinum * Tuberculinum * Ustilago * Variolinum

o

AMBRA GRISEA (Ambergris).

Mind.- (Ambra grisea)

- II Comprehension slow, has to read everything three or four times and then does not understand it.
- II Distorted images, grimaces ; diabolical faces crowd upon his fancy.
- II Embarrassed manner in company ; blushing, bashful (Fer., X-ray, Tub.).
- II Fears of becoming insane.
- II Memory impaired.
- I After business embarrassment cannot sleep, must get up and walk about (Rhus r., Sep.).
- I Difficult thinking in the morning.
- I Is not able to reflect upon anything properly, feels stupid.
- I Melancholy, sits for days weeping ; with great weakness, loss of muscular power and pain in the small of the back and constipation.

Sensorium.- (Ambra grisea)

- I Great weakness in the head with vertigo.
- I She had to lie down on account of vertigo and sensation of weakness in the stomach.
- I Vertigo with feeling of weight on the vertex ; < after sleep, < after eating.

Head.- (Ambra grisea)

- II Falling off of hair, hair turns gray early.
- II Rush of blood to head caused by music.
- I On the right side of the head, a spot, where the hair when touched pains, as if sore.

Ears And Hearing.- (Ambra grisea)

II Music aggravates the cough.

I Deafness of one ear.

I Hearing' decreases ; with cold sensation in abdomen.

I Listening to music brings on congestion to head.

I Roaring and whistling in ears, in afternoon.

Nose.- (Ambra grisea)

II Nosebleed with the menses.

I Bleeding of the nose early in the morning.

I Copious nosebleed early in bed, several successive days.

I Dried blood gathered in nose.

Teeth And Gums.- (Ambra grisea)

I Bleeding of gums.

Mouth.- (Ambra grisea)

III Ranula.

I Fetor of mouth, worse mornings.

Throat.- (Ambra grisea)

I Accumulation of grayish phlegm in throat, which is difficult to hawk up, accompanied by rawness.

I Sensation of rawness in region of velum pendulum palati.

I Sore throat after exposure to a draught of air ; stitching from the throat into right ear ; pains particularly from motion of the tongue.

Stomach. - (Ambra grisea)

I After eating, cough and gaping or gagging ; anxiety ; pressure in pit of stomach, as if the food stuck and would not go down (food descends slowly and gets into the larynx, Kali carb.) ; food is felt until it enters the stomach (Alum., Bry., Phos.).

I Eructations either empty, sour or bitter.

I Eructations with cough, so as almost to cause choking.

I Pressure in stomach and hypochondria.

Abdomen.- (Ambra grisea)

III Sensation of coldness in the abdomen.

I Coldness of one side of the abdomen (left) ; also with the deafness.

I Pain in r. hypochondriac region while lying upon it.

Stool And Anus.- (Ambra grisea)

III Frequent ineffectual desire for stool ; this makes her very anxious,

II At this time the presence of other persons becomes unbearable.

I Large flow of blood with the stool.

Urinary Organs.- (Ambra grisea)

I During urination, burning, smarting, itching and titillation in the urethra and vulva.

I Sour smelling urine.

I Urine when emitted is clouded, yellowish-brown and deposits a brownish sediment, after which the urine looks clear and yellow ; coffee ground sediment.

Male Sexual Organs.- (Ambra grisea)

I A man. aet. 80, gets asthma when attempting coition.

I Sore rawness between the thighs.

I Voluptuous itching on scrotum.

Female Sexual Organs.- (Ambra grisea)

III Discharge of blood between the periods, at every little accident, as every hard stool or after a walk a little longer than usual.

II Burning in sexual parts, with discharge of a few drops of blood.

II During urination : itching, titillation, burning of vulva and urethra.

II Menses too early and too profuse.

I In childbed obstinate constipation and tenesmus ; abdomen puffed, causing much anxiety ; so nervous she cannot attempt a stool in the presence of other people, not even the nurse.

I Menses appear seven days before time.

I Puerperal eclampsia.

I Severe itching on the pudenda, must rub the parts.

I Soreness and itching, with swelling of the labia.

I Stitches in ovarian region, when drawing in abdomen or pressing upon it.

I Thick, mucous leucorrhoea, increased from day to day, or leucorrhoea at night, of bluish-white mucus ; preceding each discharge, a stitch in the vagina.

Respiration.- (Ambra grisea)

II Asthma comes on while he is attempting coition.

II Get out of breath with the cough, which is spasmodic ; rush of blood to the head.

I Asthma of old people and children.

I Itching, scraping and soreness of the larynx and trachea.

I Oppression felt in chest and between scapula : it subsides for a short time after eating.

I Titillation in larynx with spasmodic cough.

I Titillation in throat, larynx and trachea, causing violent cough.

I Whistling in chest during breathing.

Cough.- (Ambra grisea)

III Cough in spasmodic paroxysms.

III Violent spasmodic cough, with frequent eructations and hoarseness ; with pain in region of spleen as if something were torn off.

I A kind of whooping-cough ; paroxysmal but without crowing inspiration. < after lying down.

I Choking and vomiting when hawking up phlegm from the fauces.

I Cough every evening with pain under the left ribs, as if something was torn loose there.

I Cough only at night, produced by an excessive irritation in the throat.

I Cough with emaciation.

I Deep dry cough, with accumulation of water in mouth, and subsequent rawness of the throat.

I Hollow, spasmodic, barking cough ; worse from talking or reading aloud ; with gagging after eating.

I Paroxysms of cough coming from deep in chest, excited by violent tickling in throat, evening without, morning with expectoration, generally of grayish-white, seldom of yellow mucus, of salty or sour taste ; tough grayish mucus ; worse in the morning.

I Spasmodic cough of elderly or emaciated persons.

Chest and Lungs.- (Ambra grisea)

I Itching in chest.

I Sensation of pressure deep in r. chest ; also in l. chest or in upper part of chest.

I Sensation of rawness in chest.

Heart And Pulse.- (Ambra grisea)

I Anxiety at the heart, causing oppression of breathing, with flushes of heat ; oppression beginning in the heart.

I Violent palpitation, with pressure in chest, as if a lump lay there, or as if the chest was stuffed up.

Neck And Back.- (Ambra grisea)

II Stiffness in small of back after sitting.

Upper Limbs.- (Ambra grisea)

I Coldness of left hand with headache.

I Finger nails brittle, with a very old man ; they got soft, elastic and pink.

I The arms go to sleep when lying on them, when carrying anything in the hands or at night ; with numbness especially of the left arm in rest ; tingling in the thumb.

Lower Limbs.- (Ambra grisea)

I Sore and raw between the thighs, II and in the hollow of the knees.

Nerves.- (Ambra grisea)

I Arms and limbs "go to sleep" easily.

I Great lassitude, especially mornings in bed.

I Infantile spasms.

I Jerks and twitches.

I Loss of muscular power.

I Spasms and twitches in the muscular parts.

I Weakness of the whole body ; of the knees, as if they would give way ; of the feet, with loss of sensation ; in the stomach, so that she must lie down.

Sleep.- (Ambra grisea)

II Sleepy on retiring ; but as soon as head touches the pillow is wide awake for hours, restless, tossing, or simply unable to close the eyes.

I Cannot sleep, must get up ; worryment from business embarrassment ; insomnia of business men.

I Vexatious, anxious dreams and talking in sleep ; awaken frightened.

Fever.- (Ambra grisea)

I Chill in forenoon, with lassitude and sleepiness ; relieved by eating ; skin of whole body cold, except face and genitals.

Skin.- (Ambra grisea)

II Itching.

○

ANTHRACINUM (Anthrax Poison).

Mind.- (Anthracinum)

II Anxiety, particularly in precordia.

II Delirium.

II Depression, with debility, and chill.

II Excitement.

II Loss of consciousness.

II Thinks she feels death approaching.

I Disinclined to work.

Head.- (Anthracinum)

III Headache, as if a smoke with a heating pain was passing through the head (fumée de douleur chaude) ; two shepherds who caught it from their flock.

II Confusion.

II Dizziness with pain in head.

II Headache with chill.

II Here and there in all parts of brain small and large hemorrhages of embolic origin ; after death from anthrax.

II If fully conscious they complain of great pain in head.

II Membranes of brain exhibit circumscribed or symmetrically extended bloody infiltrations.

II Pain in head, dizziness ; inner anthrax.

II Uncomfortable feeling in head, slight chills, mild fever.

I Dizziness.

I Flying gangrene, head swollen (in swine).

I Head is affected in an indescribable manner.

I Swelling of the head (sheep).

Eyes And Sight.- (Anthracinum)

II A pale yellowish or greenish swelling, if in the eyelids, of a half-translucent aspect.

II Great dilatation of pupils ; inner anthrax.

I A pale redness above the brows along the forehead.

Ears And Hearing.- (Anthracinum)

II Ringing in the ears ; inner anthrax.

I Swelling extending backward over the angle of the r. lower jaw, which could not be felt and up to near the ear.

Nose And Smell.- (Anthracinum)

- II Bloody suffusions on mucous membrane of nose.
- I Intense redness of the r. half of nose, extending to the cheek.

Face.- (Anthracinum)

- I A large stony, hard, pale swelling around the r. lower jaw, nearly painless, disfiguring the face.
- I A stony swelling around the r. lower maxilla, the inner space of the mandibula filling up to half, reaches to nearly half the cheek, and disfiguring the face, extending backwards over the angle of the lower jaw ; very little pain, not red, but sharply defined edges.
- I Beginning of swelling was in the region of the r. submaxillary gland.
- I Could not move the lower jaw as usual.
- I Could open the mouth only so far as to put the point of the tongue out.
- I Erysipelatous, dark brown redness and swelling over the whole right side of face, the nose and part of l. cheek ; swelling very hard, redness does not disappear under the pressure of finger.
- I Extending to cheek, redness from nose.
- I Gland under the chin painfully swollen.
- I Impossible to open the jaws in the least.
- I Swelling extending from the inner edge of the left lower jaw across the whole throat, in front and over the edge of the r. lower jaw, and *au niveau* with the upper surface of the r. lower molars.
- I Tearing in the right lower jaw.

Teeth And Gums.- (Anthracinum)

- I On making an incision near second molar a mass of stinking, brown ichor is discharged.

Taste, Tongue.- (Anthracinum)

- II Tongue often furred, with a thick brown coat ; dry.
- I Flabby taste.

Mouth.- (Anthracinum)

- II Bloody suffusions and hemorrhagic collections on the mucous membranes of canthi of mouth and nose ; inner anthrax.
- II Continued bleeding from the mouth ; the blood shows a lack of power to coagulate ; with inner anthrax.
- II Dark red, bloody ecchymoses of mouth.
- II Superficially escharred pustules in mouth after death.
- I Fundus of mouth is elevated by the swelling, as hard as a callus, extending back to the parotids, and reaching up to the external surface of the lower jaw.

I Offensive odor from mouth. Mouth could not be opened.

I Saliva increased.

Throat.- (Anthracinum)

II Slight difficulty in swallowing ; inner anthrax.

II The submucous tissue, especially in fauces and around the larynx, is thickened and oedematous.

I Could not swallow, with great thirst.

I Cynanche cellularis ; a sharply-marked margin about the swellings.

I Region of the throat above the larynx to the mouth swollen.

I Right tonsil hurts.

I Swallowing exceedingly difficult.

Appetite.- (Anthracinum)

III Loss of appetite, prominent in every patient.

II Loss of appetite and gastralgia ; inner anthrax.

II Loss of appetite, with chills.

I Excessive thirst, but can hardly swallow.

I Symptoms from putrid water.

Stomach.- (Anthracinum)

II Gastralgia.

II Mucous membrane of stomach and intestines reddish, swollen, with isolated or numerous oedematous, hemorrhagic prominent infiltrations, from size of a lentil to that of a coffee bean, showing a grayish or greenish-yellow discolored surface, with a positively sloughing centre.

II Nausea and vomiting with chill.

II Numerous peculiar hemorrhagic and superficially eschared infiltrations of stomach and intestines ; intestinal anthrax.

II Vomiting followed by diarrhoea.

II Walls of stomach and intestine oedematous, discolored, a cloudy red.

I Pressure and burning in the region of the stomach.

Abdomen.- (Anthracinum)

II Bellyache with chill.

II Colicky pains ; inner anthrax.

- || Enlargement of spleen.
- || Epidemic spleen disease of cattle or horses.
- || In the intestines a thin fluid material, slightly colored with blood.
- || Mesenteric and retro-peritoneal glands enlarged to the size of a walnut ; form blackish-red masses, held together by a jelly-like congestive tissue, infiltrated with serum.
- || Moderate serous or sero-hemorrhagic effusion and sub peritoneal suggilations.
- || Mycosis intestinalis ; intestinal anthrax.
- || Peculiar pustular and carbuncular foci in the intestinal tract.
- || Serous and sero-hemorrhagic infiltrations of the peritoneal and mesenteric connective tissue, walls of stomach and intestine, and of the mucous membranes.
- || Simple hemorrhages, infarctions and foci on different parts of intestines.
- || Swelling of the abdomen ; inner anthrax.
- || The retro-peritoneal and mesenteric connective tissue infiltrated, jelly-like, and of a yellowish-reddish color.
- || The same disease in sheep. *Anthracinum suum* is better than *Anthracinum ovium* in the acute form, but in the chronic form *Anthracinum ovium* is better.
- I Sensation as if the diaphragm was pushed forward.
- I Sensation of anxiety and constriction, most in the praecordia, liver engorged, slight hemorrhage here and there, spleen moderately enlarged, soft, full of blood, dark color.

Rectum And Stool.- (Anthracinum)

- || Diarrhoea.
- || Vomiting followed by a painless, moderate, more or less intense, often bloody diarrhoea ; inner anthrax.
- || With the diarrhoea sometimes a cholera-like collapse ; inner anthrax.

Urinary Organs.- (Anthracinum)

- || Kidneys swollen, with oedema, sprinkled with small hemorrhages, engorged ; suggilations in mucous membranes of the pelvis.

Respiration.- (Anthracinum)

- || Breathing frequent, laborious ; quick, spasmodic ; inner anthrax.

Lungs.- (Anthracinum)

- || Oedema of the mediastinal lymphatic glands.
- || Slight serous effusions into pleural cavities.
- || Sub-pleural ecchymoses with vascular engorgement, and a dark coloring of the parenchyma.

Heart, Pulse And Circulation.- (Anthracinum)

- II Blood not coagulating.
- II Blood of a dark cherry red, generally fluid or with some loose clots.
- II Cyanosis ; inner anthrax.
- II Discolored lines over the veins, or red lines and stripes in the course of the lymphatics.
- II The beating of the heart stronger, more decided and more perceptible.

Neck And Back.- (Anthracinum)

II Carbuncle on the back, nine inches in length and five inches in its greatest width ; with sloughing, abundant discharge of ichorous, terrible smelling pus, and blood-poisoning by absorption of pus.

Upper Limbs.- (Anthracinum)

- II Felon, the worst cases, with sloughing ; severe torturing pain and great prostration.
- I Arms and hands covered with a crusty eruption, full of cracks, discharging pus and an acrid fluid, with painful, unbearable itching ; checked for a while by the Old School, it had burst out again with terrible fury.
- I On the middle of the palm of the hand a large blister, which, when opened, discharged a yellow watery fluid.
- I The whole left hand (not the fingers) swollen, highly reddened, very painful ; the redness extended over the whole hand and even the wrist, and a red streak ran up the forearm.
- I Whitlow.

Lower Limbs.- (Anthracinum)

- II Severe pains in limbs and joints with the fever ; intestinal anthrax.
- I Above the knee, redness, swelling and pain, and later a large black blister on inside of thigh, extending four inches upward and inward ; after being lanced bloody water ran out.
- I Bluish-brown spots which break open.
- I Foot oedematous.
- I From the openings on lower leg, caused by the fracture, a copious stinking pus (like carious bones).
- I On outside of knee a large fluctuating swelling, by pressure discharging a horribly smelling gangrenous ichor.
- I The whole thigh was swollen, most above the knee, and also the foot.
- I Thighs livid to the nates, hard and painful ; lower legs dark blue, feet oedematous ; when the blisters break they discharge an offensive ichor.

Nerves.- (Anthracinum)

- II Cholera-like collapse after diarrhoea.
- II Clonic spasms, trismus or opisthotonos ; sometimes in serious cases.

II Debility and depression, with pains in limbs and general sense of malaise, followed by disturbance of intestinal canal ; inner anthrax.

II Debility with chill.

II Epileptiform convulsions ; inner anthrax.

II Great restlessness.

II Great weakness with the fever.

II Marked rigor mortis after death.

II Opisthotonos ; inner anthrax.

II Sudden fatal issue, preceded by extreme collapse.

II Tetanic spasms in upper limbs.

II With cyanosis, asphyxia and the most extreme collapse, followed by death in all cases of bleeding.

I Completely exhausted, she thinks she feels death.

Sleep.- (Anthracinum)

II Restless sleep, with chill.

II Somnolence ; inner anthrax.

I Restless, irritated at night.

I Sleeplessness.

Fever.- (Anthracinum)

II Chilly, with debility, headache, followed by a general malaise, loss of appetite, restless sleep, great debility and depression, and in eight or ten days carbuncles, most on the arm, forearm, head.

II Cold sweats in serious cases.

II Decided chill, followed by bellyache, nausea, vomiting and in two or three days with the supervention of collapse and cyanosis ; death.

II Febrile movement, slight in the beginning, is often followed apace by high fever ; great weakness, delirium, excitement, confusion.

II Slight chills with fever and strange sensation in head.

II Sweat all over with debility.

II Temperature very slightly elevated ; inner anthrax.

I Copious sweat.

I Disposed to sweat ; rather sticky.

I Fever with diarrhoea.

I Heat, thirst, less appetite, suffering and fatigued.

I Moderate heat, little thirst, general sweat.

I Very much fever.

Skin.- (Anthracinum)

III Anthrax carbuncles cured by Anthracine, every day, also externally, in four days.

III Carbuncle with horrible burning pains ; or discharge of ichorous offensive pus.

II A little red speck, like a flea-bite, with a central black point, swells gradually and changes into an itching papule, capped with a small, clear, reddish or bluish vesicle, gradually enlarging.

II Anthrax pustules most on face, forearms, hands, fingers, neck, less often the ear, still less frequent the covered parts.

II Black blisters, fatal in twenty-four to forty-eight hours.

II Carbuncle darkish red, greasy, and is often more eroded than ulcerated.

II Carbuncle on arm, forearm, head. See chill.

II In case of more than one detritus, the whole is swollen like erysipelas, and when cut it looks like the *Vespajas* of the Italian dermatologists.

II Large cutaneous eschars.

II Little carbuncles ; inner anthrax.

II Papules and pustules, with extensive oedematous and phlegmonous infiltration of the neighboring skin and subcutaneous tissue.

II Small and large epidermal vesicles filled with serum.

II The anthrax pustule penetrates deeply into the subcutaneous cellular tissue.

II The dense or doughy soft papules or pustules, around and beneath the eschar, vary in size from a pea to a nut.

II The excoriated spot dries up, becomes brown and livid and a local eschar forms.

II The papule promptly bursts and discloses a dark red base.

II The secondary vesicles contain a yellowish, reddish and blackish fluid.

II With an increase of the round thick eschar, one-fourth to three-fourths of an inch, the raised border also extends.

I After having taken homoeopathic medicine for malignant ulcers, suddenly the greatest malaise, and a black blister formed below the knee with swelling all around, and feverish shaking chill through the whole body.

I All openings run into one, discharge much pus.

I Black or blue blisters.

- I Blister on palm of hand.
- I Carbuncle.
- I Chronic forms of anthrax with indurations like knots under the skin.
- I Crusty eruption discharging acrid fluid.
- I Crusty oozing eruption, with the most violent itching.
- I Erysipelas gangrenosa.
- I Erysipelatous form of chronic anthrax.
- I Erysipelatous inflammation about the carbuncle.
- I Itching with dry skin ; violent as if mad (horses).
- I Large black blister on inside of thigh.
- I Most malignant gangrenous ulcers (sheep).
- I Seventh day after the remedy several larger and smaller openings, discharging watery, sometimes bloody matter, very little pus ; swelling less hard around the base.
- I Skin dry, itching violently and burning.
- I Small-pox of sheep.
- I Ulcus excedens (sheep).
- I Unbearable itching on arms and hands.

- o **ELECTRICITY.**

Mind.- (Electricity)

- II Involuntary hysterical laughter.
- II Loss of memory.
- II Weeping, timid, fearful ; sighing, crying out through nervous fear.

Sensorium.- (Electricity)

- II Loss of sensibility.
- II Stupefaction.

Head.- (Electricity)

- II Headache ; pressure in the forehead, from above downwards, as from a stone.
- I Tearing from the nape of the neck to the forehead.

Eyes.- (Electricity)

II Blindness.

II Everything looks yellow.

II Inflammation of the eyes ; profuse lachrymation.

I Black point before the right eye.

Ears.- (Electricity)

II Blisters behind the ears full of an acrid fluid.

Nose.- (Electricity)

I Loss of smell.

Face.- (Electricity)

I Large blisters on the cheeks.

Mouth and Pharynx.- (Electricity)

II Foam at the mouth.

II Increased secretion of saliva.

II Swelling of the tongue.

II The tongue is very sensitive, particularly at the tip.

Stomach, Nausea and Vomiting.- (Electricity)

II Heartburn.

Abdomen.- (Electricity)

II Cutting in the abdomen at the approach of a thunder-storm.

Urinary Organs.- (Electricity)

II Frequent micturition.

I Sensation as if the bladder would burst.

Female Sexual Organs.- (Electricity)

I Black and thick menstrual blood.

Back.- (Electricity)

I Boils in the back and nape of the neck.

Upper Limbs.- (Electricity)

|| Feeling of numbness in the tips of the fingers.

|| Frightful pains in the arms and lower limbs.

|| Paralysis of the arms.

| Swelling of the hand, also red, or sudden.

| Trembling of the hands.

Lower Limbs.- (Electricity)

|| Burning in the feet, up to the knees, particularly at night.

Skin.- (Electricity)

|| Ecchymoses.

|| The skin becomes blackish.

General Symptoms.- (Electricity)

|| Drawing through all the limbs, extending to the tips of the fingers and toes.

|| Fainting.

|| General languor after a meal.

|| Pains in the limbs.

|| Paralysis of single limbs, particularly the lower.

|| Stiffness of the limbs.

|| Trembling of the limbs, particularly of those which have received the shock.

○ LAC CANINUM.

Mind.- (Lac caninum)

|| After inhaling gas for extraction of teeth, very strange sensation in head (such as he felt when going off under gas) ; sometimes imagines heart or breathing is going to stop, or otherwise frightens himself, and this makes heart beat violently ; occasionally very depressed, and fancies he is going out of his mind.

|| Attacks of rage, cursing and swearing at slightest provocation.

|| Cannot bear to be left alone for an instant.

|| Child cries and screams all the time, especially at night, and will not be pacified in any way.

|| Chronic " blue " condition ; everything seems so dark that it can grow no darker.

|| Easily excited.

|| Exceedingly nervous and irritable.

|| Fear of death, with anxious expression of countenance.

|| Fears she will become unable to perform her duties.

|| Feels weak, and nerves so thoroughly out of order that she cannot bear one finger to touch another.

|| Gloomy feelings, < as headache gets worse.

|| Imagines that he wears some one else's nose.

|| Imagines to be dirty.

|| No desire to live.

|| Thinks that she is looked down upon by every one, that she is of no importance in life, and feels insulted thereat.

|| Too excited to allow examination of throat.

|| Very easily startled.

|| Very nervous ; constant dread ; a feeling as if she was going to become unconscious.

|| Wakes distressed, and obliged to rise and occupy herself in some manner ; fears she will be crazy.

|| When awake, very irritable and cries constantly.

|| When paroxysms of intense nervousness come on, feels like tearing off her clothes ; takes off her rings ; cannot bear anything to touch her, especially over l. ovarian region, from which she frequently lifts bed clothes.

| Cannot collect her thoughts ; confused feeling.

| Depression of spirits, doubts her ability and success, thinks she will have heart disease and die of it.

| Imagines she sees spiders.

| Imagines she sees spiders.

| Intense ugliness and hatefulness ; writes to her best friends all sorts of mean and contemptible things.

| Omits final letter or letters of a word, when writing ; in speaking substitutes name of objects seen, instead of object thought of.

| Sensation as if she were going deranged, when sitting still and thinking ; sometimes she has most horrible sights presented by her mental vision (not always snakes), feels horribly afraid that they will take objective form and show themselves to her natural eye.

| Very absentminded ; makes purchases and walks off with out them ; goes to post a letter, brings it home in her hand.

| Very forgetful ; in writing, uses too many words or not the right ones ; very nervous.

Head.- (Lac caninum)

- III After midnight, very severe frontal headache, and a piercing pain on vertex.
 - III Headache both frontal and occipital, < by turning eyeballs upward.
 - II After inhaling diphtheritic breath, light headed, with tingling on vertex and slight sore throat.
 - II Constant noise in head, very confusing ; < at night and at menses.
 - II Darting pains from occiput to forehead ; beating.
 - II Frequently wakes with sick headache, which seems to commence at nape.
 - II Headache from below eyes over whole head and top of shoulders.
 - II Headache over eyes, < when sewing.
 - II Headaches mostly through temples, darting, stabbing ; sometimes begin on right side and sometimes on left ; always going from one side to the other.
 - II Headaches seem unbearable, and are attended by pain in lumbar region ; all pains cease as soon as throat gets <.
 - II Pains and throbbing in temples.
 - II Severe pressure on brain.
 - II Sick headaches, beginning in nape ; pain settling gradually in right or left forehead.
 - II Wakes at night with sensation as if bed was in motion ; noise in head bad beyond description ; first thought on waking that headboard was swaying, and so occasioning distress, but found it arose from internal causes.
 - I Arose in morning with heavy, dull, frontal headache, and at 9 A. M., severe sharp pain on top of head, coming from nape of neck, then stretching across head forward ; pain so severe that he presses top of head with hands ; neck stiffened ; bending head forward, or lying clown, causes congestion, increasing pain ; again pains subside for short time, and begin anew, either in front part of head, or in nape of neck, or all over head at once ; when pain is frontal it causes lachrymation.
 - I Dizzy sensation with slight nausea.
 - I Frontal headache.
 - I Headache : in afternoon, principally over left eye ; over left eye on first awaking.
 - I Headache < by noise or talking, > by keeping quiet ; confused feeling in head.
 - I On going into cold wind, terrible pain in forehead as if it would split open, > on going into warm room.
 - I Pains in head during day, first on one side, then on other seem perfectly unbearable ; > on first going into air, but soon grow
- Eyes.- (Lac caninum)**
- II Agglutination of left eyelids.
 - II Burning in left eye.
 - II Dark brown areolae under eyes.

- II Difficulty in distinguishing objects ; in reading, letters run together.
- II Eyes blurred.
- II Eyes dull and lustreless.
- II Eyes sensitive to light.
- II Eyes watery and discharging.
- II Film over eyes when reading or looking closely.
- II Frequent sensation of a film before eyes, with vertigo, and while thus suffering would see a small dark object, like a mouse or bird, coming up to her left.
- II Heaviness of left upper lid, with pain above left eye.
- II Heavy pain in the eye-balls with outward pressure.
- II Must have light, yet is intolerant of sunlight.
- II Non-oedematous swelling of upper and lower lids ; pink color of lower lids ; most noticeable on right.
- II Sees big eyes and creeping things.
- II Sees faces before her eyes, < in the dark ; the face that haunts her most is one that she has really seen.
- II Square or round green spots or brown spots before left eye, when sun is bright ; sometimes bright spots before left eye.
- II When looking in a mirror by gaslight, after exerting eyes, sees a green spot or a green band before her left eye, the band slanting downward from left eye to r. cheek.
- I Tendency in retina to retain the impression of objects, esp. of colors ; or somewhat of the object last looked at is projected into the next.
- I Upper eyelids very heavy, could scarcely keep eyes open.
- I When reading the page does not look clear, but seems covered with various pale spots of red, yellow, green and other colors.

Ears.- (Lac caninum)

- II Green, odorless discharge.
- II More than any other remedy, relieved deafness from hereditary syphilis.
- II Sounds seem very far off.

Nose.- (Lac caninum)

- III Coryza, with discharge of thick white mucus.
- III One side of nose stuffed up, the other free and discharging thin mucus at times and thin blood ; these conditions alternate, first one nostril stopped up and the other fluent, and *vice versa*.
- III Profuse nocturnal nasal discharge, like gonorrhoea, staining pillow greenish yellow.

II Epistaxis : when speaking or swallowing ; at 4 P. M., returning at intervals.

II Fluids escape through nose while drinking.

II Nose became so bad that there was fear of destruction of bones ; bloody pus discharged several times daily ; nasal bones sore on pressure.

II Nose cold.

II Two very angry gatherings, one under left side of nose, and one on upper left nostril ; both came to a head, and discharged matter and blood, and afterwards scabbed over ; before discharge shooting pain.

I Considerable sneezing.

I Coryza ; constant watery discharge from nose, excoriating nostrils and upper lip.

I Fluent catarrh from both nostrils, with sensation of fullness in upper part of nose.

I Head so stuffed she can hardly breathe.

I Nasal discharge, excoriating nostrils and upper lip.

I Sore on right side of septum of nose ; next day nose sore, constant inclination to pick at it and get the scab off ; nose still sore on sixth day, and on seventh day was very painful to touch ; but on eighth day scab came off nose, leaping it as well as ever.

I Stuffed feeling in head, as of a severe cold in head.

Face.- (Lac caninum)

II Countenance pale and careworn.

II Face flushed, swollen and hot ; burns, feels dry.

II Flushes on left cheek.

II In morning, left superior maxillary feels sore ; most of time there is dull pain, < by exertion ; sometimes throbbing pain, burning, throbbing, aching heat, sensation of fullness ; cannot wear her false teeth from soreness and swelling of maxillary ; an exacerbation of pain leaves face very sore ; pains > by warm applications, but only cold applications > soreness.

II Parotid gland first attacked, and disease extends to other glands of neck ; throat and sides of neck not tender to external touch.

II Seems to affect lower lip most, and blisters and fever sores on lip are amenable to its influence.

II Submaxillary glands swollen.

II Swelling of left parotid, with sore throat and loss of appetite.

II Swelling of parotid passes from right to left but more often from left to right.

I Dark brown areola under eyes.

I Face flushed ; cheeks red.

I Lips dry and peeling off ; dry and parched, but mouth constantly full of frothy saliva.

Teeth.- (Lac caninum)

- II Gums swollen, ulcerated, retracted, bleeding, teeth loose ; caused by defective nutrition and exposure.
- II Paroxysmal gnawing pain in left upper canine, temporarily yields to any cold application.

Mouth.- (Lac caninum)

- II Nothing tastes natural, except salt food.
- II Slight yellow coating on tongue.
- II Breath offensive, putrid.
- II Constant spitting and drooling, very profuse, making chin and breast sore.
- II Difficulty in articulating, owing to a semi-parietic state of tongue, causing stuttering if she talks fast ; has to speak very slowly.
- II Mouth dry and parched ; drinks little and often.
- II Peculiar rattle in mouth, right along tongue ; on attempting to hawk her mouth clear, mucus rattled along tongue, quickly and continually ; utterance was so indistinct as to be unintelligible, and every word she tried to speak was accompanied by this quick and continuous rattle along tongue.
- II Stomatitis ; stomacace ; cancrum oris ; nursing sore mouth.
- II Tongue : heavily coated, and dry to the tip ; dirty coated, yellowish-white and slimy ; dry ; thickly coated, grayish-white.
- I During sleep saliva runs from mouth so as to wet. pillow.
- I Frothy mucus in mouth < by going into open air, and after eating.
- I Increase in quantity of saliva which is slightly viscid.
- I Mouth constantly full of frothy mucus, but a constant inclination to swallow.
- I Mouth constantly full of frothy saliva, lips dry, parched.
- I Mouth very dry, without thirst.
- I Tongue coated brown.
- I Tongue coated whitish grey, having an underlying bluish look.
- I Tongue coated whitish, except edges, which are red.
- I Tongue coated, dirty looking, centre to root.
- I Tongue furred, whitish edges, centre and root darker.

Throat.- (Lac caninum)

- III Diphtheria and diphtheritic croup ; membranous croup.

III Diphtheritic membrane white like china ; mucous membrane of throat glistening as if varnished ; membranes leave one side and go to the other repeatedly ; desire for warm drinks, which may return through nose ; post-diphtheritic paralysis.

III Soreness of throat commences with a tickling sensation, which causes constant cough ; then a sensation of a lump on one side, causing constant deglutition ; this condition entirely ceases, only to commence on the opposite side, and often alternates, again returning to its first condition ; these sore throats are very apt to begin and end with menses.

III Swallowing very difficult, painful, almost impossible ; fluids return through the nose.

III Tonsils inflamed and very sore, red and shining, almost closing throat ; dryness of fauces and throat ; swelling of submaxillary glands.

II After membrane exfoliates, mucous membrane appears raw and bloody, with increased deglutition.

II Both tonsils swollen and covered with spots of exudation, like the mould on preserves.

II Dark red, angry streaks of capillaries in fauces, giving place to shining, glistening deposit, or tough membrane ; half arch is filled with sticky, fetid saliva.

II Dark red, angry streaks of capillaries in fauces, giving place to shining, glistening deposit, or tough membrane ; half arch is filled with sticky, fetid saliva.

II Diphtheritic deposits look as if varnished ; exudations migratory, now here, now there. constantly reforming ; swelling in throat so large and tense that mouth could not be closed.

II Diphtheritic deposits look as if varnished ; exudations migratory, now here, now there.

II False membrane in throat, thick, grey, or slightly yellow, or dark and almost black, or white and glistening, almost like mother of pearl, or fish scales.

II False membrane, thick grey, yellow or dark, surrounding mucous membrane dark or bright, may be < on either side, or inflammation shift from side to side, generally < on left

II False membrane, thick, yellowish grey, often greenish.

II Gargling with warm water brought up a stringy mucus.

II In most cases of diphtheria, the throat symptoms begin on right side.

II Inflammation, ulcers and swelling shift from side to side, generally < on left

II Lump on left side of pharynx below tonsil, causing an enlargement that filled each arch of palate, nearly to right side.

II Marked sensation of lump in throat on left side, when swallowing ; pain extends to ear.

II Membrane would leave throat, and a very severe interstitial hemorrhage of bright red blood would ensue ; hemorrhage would slowly improve, and membrane appear again in throat ; these had continued to alternate for several days.

II Most pain when swallowing solids, no aversion to cold drink ; when swallowing solid food it seems to pass over a lump, with sore and aching pains extending to and into left ear.

II Mucous follicles raised or swollen, and covered with a whitish, cream colored mucus.

- || On each tonsil a very thick exudation, covering nearly entire surface ; while examining a large piece of membrane was accidentally detached from one tonsil, followed by considerable hemorrhage.
- || On each tonsil a very thick exudation, covering nearly entire surface ; while examining a large piece of membrane was accidentally detached from one tonsil, followed by considerable hemorrhage.
- || On swallowing acute pain at one time on right side of throat, and again on left side.
- || Paralytic symptoms strongly marked ; as soon as he went to sleep would stop breathing, and was only kept alive by keeping him awake ; apparently respiration was kept up by voluntary effort.
- || Pharyngeal inflammation, with wholesale destruction of epithelium, viscosity of saliva, heat of palms ; absolute necessity for constant change of position.
- || Quinsy : suppuration ran from left tonsil to right, then from right to left, then back again to right, then both tonsils equally ; and again one tonsil would > and the other grow < ; whole posterior portion of throat was an oedematous swelling, rising up like an insurmountable barrier ; thick tough pieces of diphtheritic membrane were coming away, and new membranes constantly reforming ; swelling in throat so large and tense that mouth could not be closed.
- || Right tonsil raw, swollen ; grey white membrane there and on fauces.
- || Sensation as if throat were closing and she would choke, sensation is between throat and nose ; feels as if something in throat was either enlarged or relaxed ; desires to keep mouth open lest she should choke ; sometimes cannot swallow, because there seems to be a kind of muscular contraction in throat.
- || Sensation of ball or round body in left side of throat, and feeling that it could be removed with a knife.
- || Sensation of lump in right side of throat, with a feeling that she could take hold of it with her fingers and pull it out ; accompanied by a very annoying pricking, stitching feeling ; constant inclination to swallow saliva, which causes soreness of throat.
- || Sensation of rawness, commencing usually on left side of throat.
- || Small, round or irregular, grey white ulcers, on tonsils and fauces, both sides.
- || Sore spot on left side of throat, only at night, removed by 1 A. M. ; next night same on right side of throat ; after 1 A. M. returned no more.
- || Sore spot on left side of throat, only at night, removed by 1 A. M. ; next night same on right side of throat ; after 1 A. M. returned no more.
- || Sore throat just before menses for several years ever since diphtheria ; small yellowish white patches of exudation on tonsil of affected side, with great difficulty of swallowing, and sharp pains moving up into ear ; these patches are also present on back of throat and uvula ; some are quite yellow and some are white ; scraping them off makes them bleed.
- || Sore throat on right side, low down, and extending up to ear ; pain when swallowing ; sensitive to external pressure ; slight coryza.
- || Sore throat, pains in whole body and limbs, severe headache.
- || Sore throat, ulcer on inner side of each tonsil, tonsils red and slightly enlarged, rest of the throat dry.
- || Talking is very difficult and there is a disposition to talk through nose.
- || Thick membranous mass lying on soft palate, left side ; diphtheritic masses covering uvula and posterior wall of throat ; next day, membrane on soft palate thicker, dirty brown on uvula and posterior walls and pillars

of throat, much more extensive and offensive ; very difficult deglutition ; a large membranous mass, which threatened suffocation, having been removed by forceps ; on following morning a second membrane had taken place of first, and walls of throat were covered with a dirty grey exudation ; uvula almost black, and coarse shreds of membrane hanging from it.

|| Thick, dirty grey diphtheritic membrane covering free border of epiglottis, and extending off to each side.

|| Throat > after drinking cold or warm ; < by empty swallowing.

|| Throat covered with diphtheritic membrane ; uvula elongated, swollen and covered with black and white or grey diphtheritic deposit ; back of throat extending to hard palate, all covered ; breath very offensive ; left side of neck swollen and almost even with jaw ; great difficulty in swallowing ; after throat began to improve, disease seemed to work through whole alimentary canal, for uvula and parts were very much swollen, and every little while there would be involuntary discharges of diphtheritic matter from uvula and rectum.

|| Throat filled with substance that looked like "smear käse ; " throat, tongue, roof of mouth, gums and cheeks completely lined with this substance ; mouth and throat filled with loose particles ; horrible odor.

|| Throat has a burnt and drawn feeling as from caustic.

|| Throat highly inflamed, swollen, almost closed, grey diphtheritic spots on left side of throat.

|| Throat sore, but little swelling, tonsils very slightly enlarged ; soreness of throat, first chiefly right then left ; well marked diphtheritic membrane on both sides of the throat situated on an inflamed red base, 1/2 inch long, 1/2 inch wide, 3/8 inch thick, and the same length and width as at the base ; anterior edge a dirty yellow ; centre more organized, pearly, glistening, white like cartilage ; membrane on right side seems more firm and dense, and disappeared later.

|| Throat sore, oedematous, puffed, tonsils badly swollen.

|| Throat sore, swollen, red and glistening.

|| Throat very dry and sore, much inflamed, < right side ; palate red, uvula elongated ; very painful deglutition.

|| Throat very sensitive to touché externally.

|| Throat very sore, < left side ; large greenish ulcers on both tonsils, surrounded by grey white exudation, parts not covered are a deep purple red ; swelling externally on both sides ; after exudation on tonsils disappeared, a raw, bloody surface was left.

|| Throat very sore, tonsils enlarged, especially left, very large white patches ; tonsils and pharynx deep purple red ; putrid odor from throat ; after patches were expectorated, they left throat **very** sore, raw and bloody.

|| Tongue, fauces, tonsils, all swollen and covered with a dirty coating.

|| Tonsils swollen as almost to close throat.

|| Ulcers increase in size and number, but neighboring membrane looks clearer.

|| Ulcers on throat shine like silver gloss, symptoms went from side to side ; croupy symptoms not well marked ; after exudation was cleared off, a deep excavation was left.

|| Uvula elongated and very much swollen, diphtheritic coating on it ; tonsils swollen and coated ; back of throat patched, extending up to hard palate ; odor offensive and diphtheritic.

|| Uvula pretty free from membrane, but intensely sore and bleeds.

II Wakes with throat and mouth painfully dry.

II White patches, like eggs of flies, on both tonsils, extending thence to back of throat ; tonsils enlarged and deep red ; felt she would suffocate at night from full feeling in throat, which prevented sleep ; swallowing toast gave some pain, but seemed to clear throat ; and she had to gulp it down.

II White ulcers on tonsils.

II Whole membrane of throat highly inflamed, swollen, and glands enlarged on both sides.

II Whole membrane of throat swollen, dark red, with grey patches and small, irregular shaped ulcers ; membrane peeled off occasionally ; articulation and deglutition intensely painful ; < after sleep.

II Whole of right tonsil covered with diphtheritic patch.

I Constant inclination to swallow, causes pain extending to both ears.

I Crusts on skin, with grayish yellow matter under them ; mucous follicles of throat raised and swollen, and covered with whitish, cream colored mucus ; bloody pus discharged from nose several times a day ; nasal bones sore to pressure.

I Fever : bathed in warm perspiration, especially about the face, neck and hands ; anxious expression ; eyes watery and discharging ; wants to sit up in mother's arms ; cries and desists at every attempt to nurse ; reaches for water, yet refuses to take it ; respiration hoarse ; crying whispered and broken, often no sound at all ; pulse **170 ; tongue, fauces and tonsils swollen and covered with dirty coating ; drooling from mouth ; throat tender to touch externally ; thick, dirty, grey, diphtheritic membrane, covering free border of epiglottis, and extending off to each side ; child refuses to swallow and sputters out the medicine, some returning through the nose.**

I Glossy, shining appearance ; disposition on part of membrane to change its position in fauces.

I Heaviness, and stomach bloated and tender ; enlargement of tonsils, left tonsil < ; feels weak ; cannot eat or drink anything without pain in pit of stomach ; shortness of breath and general languor.

I In morning throat very sore ; right tonsil covered with ulcers and patches, which extended over palate and covered left tonsil ; next day membrane extended across posterior wall of pharynx ; uvula elongated, accompanied by chilliness, high fever, pains in head, back and limbs, great restlessness and extreme prostration. This was pronounced to be "severe diphtheria." but it soon got well.

I On third day right tonsil swollen and on it a small diphtheritic patch, rest. of throat inflamed ; on fourth day both tonsils swollen and covered with diphtheritic patches, with difficult deglutition ; high fever, restlessness, cried out and talked in sleep ; complained of pains in head, back and limbs ; bright scarlet redness on chest and around neck, which, on fifth day, extended all over body and legs ; disease now at highest point ; skin, in large patches, assumed a dark red color bordering on purple ; whole body swollen ; membrane, swelling and soreness on right side ; deglutition impossible ; refusing to drink while complaining of intense thirst ; characteristic fetor in room ; soreness on right side decreased and commenced on left ; left tonsil and posterior wall of pharynx covered with membrane ; posterior nares invaded ; marked sensation of lump in throat on left side, when swallowing, with pain extending to left ear. ; tongue coated dirty, yellow white and slimy ; absence of prostration ; improvement commenced on seventh day and remedy was discontinued.

I On waking in morning, throat felt as if there were lumps in it like two eggs, and sore all the time, especially when swallowing ; cold water seemed to > momentarily ; in evening, examination revealed both tonsils much swollen and very red, left most, and distinct patches on left tonsil.

I Pain in throat pushes toward left ear.

I Pains in limbs, small of back and head disappear, and throat becomes more painful, but looks better ; often ulcers increase in size and number, but neighboring membrane looks clearer ; < by empty deglutition ; throat

feels stiff ; > after drinking, warm or cold, no thirst, but dry mouth ; pain pushes toward left ear ; right tonsil raw, swollen, grey white membrane there and on fauces ; epistaxis when speaking or swallowing, in one case ; sweat all over ; great exhaustion with poisoned feeling ; frequent micturition, urine dark ; restless, legs and whole body ; face burns dry ; constant spitting, drooling ; ulcers small, round or irregular, grey white ; voice hoarse, interrupted by weakness and hoarseness.

I Partial suppression of urine ; throat sore and of an oedematous, puffy appearance ; next morning, pulse 130 ; temperature 102° ; tonsils badly swollen ; great indisposition to take food or drink.

I Patch of diphtheritic membrane appeared first on right tonsil, then on left, and frequently alternated sides ; swelling of neck (submaxillary and lymphatic glands) also alternated in like manner ; < during and after a cold storm from northeast ; tickling in throat when drinking ; one side of nose stopped up, the other free and discharging thin mucus at times and thin blood ; this condition of nose also alternated ; non-oedematous swelling of eyelids, pink color of under lid, particularly of r' eye ; breathing hoarse and croupy, at times entire stoppage of breath ; often snoring, and only possible through mouth ; obstinate constipation, frequent desire, with darting pains in rectum, no power to expel ; stool large in size, whitish, rough, scaly, hard ; could not bear to be left alone an instant ; saw big eyes and creeping things ; must have light, yet is intolerant to light of sun ; urine scanty, infrequent, no desire, coffee colored ; 80 per cent. of albumen and much mucus ; quantity less than a gill in twenty-four hours.

I Pricking sensation in throat, as if full of sticks.

I Quinsy just ready to discharge, disappeared without discharging, in an unusually short time ; the trouble had been changing from one side to other and back again ; has not returned.

I Right tonsil red and swollen ; pain in tonsil of gnawing character ; < at night ; dreams of snake in bed.

I Roof of mouth and back wall of pharynx coated with a greyish yellow deposit, greater part of which soon disappeared, lasting only about an hour ; throat very much > by noon, deposit had nearly disappeared, but < again by night.

I Severe chills, headache, pain in back and limbs, restlessness and sore throat ; three days later right tonsil covered with ashy grey membrane, extending along free palatine border to uvula, which it had already involved ; peculiar diphtheritic odor in room ; pulse small ; skin clammy ; rapid vital exhaustion ; next day membrane involved whole arch of palate and passed down to left tonsil.

I Shortly before going to bed, throat began to feel raw and sore ; did not sleep well ; next morning, throat felt full and sore, somewhat < on right side ; this condition continued two days, when it seemed to continue downward to chest.

I Shortly before going to bed, throat began to feel raw and sore ; did not sleep well ; next morning throat felt full and sore, somewhat < on right side ; this condition continued two days, when it seemed to continue downward to chest.

I Sore throat on left side ; painful to external pressure both sides.

I Sore throat, rapidly growing < ; fever ; difficult swallowing ; right side < ; right tonsil intensely inflamed, bright red and greatly enlarged, and a yellowish grey spot on inner surface ; whole pharynx, uvula and velum much inflamed ; spot became larger, and others formed in pharynx ; left tonsil became nearly as large as right ; fetid breath ; subsequently a bright scarlet eruption on face, neck, hands and chest, like scarlatina ; almost total inability to swallow, especially fluids ; aversion to liquids, particularly water.

I Soreness of throat commences with a tickling.

I Soreness of throat, accompanied by intense heat ; pulse scarcely to be counted ; prostration so complete that patient refused even to make an effort to take medicine ; temperature 102.6° ; great sensitiveness of throat externally ; symptoms after sleep ; very thick exudate, covering nearly entire surface of each tonsil, which, if forcibly removed, is followed by considerable hemorrhage.

I Throat feels raw.

I Throat highly inflamed, swollen, almost closed ; grey, diphtheritic patches on left side of throat ; difficult breathing, at times suffocative spells, pulse 140 ; face flushed, swollen and hot ; tongue dry and thickly coated, greyish white.

I Throat sore ; with severe headache ; pain extending to chest ; dry and sore ; deep red color on either side of throat opposite tonsils ; on left side ; painful to external pressure on both sides.

I Uvula coated (in seven cases).

Appetite.- (Lac caninum)

II Appetite and strength failing ; dislike to food, especially fat or greasy.

II Aversion to liquids, especially water.

II Craves milk and drinks much of it.

II Great thirst for large quantities, often.

II Thirst for a little at a time, but often, as throat is so dry and hot.

II Thirst.

I Desire for highly-seasoned dishes, which is very unusual ; has used pepper, mustard, and salt freely.

I No appetite.

Stomach.- (Lac caninum)

III Great faintness of stomach and nausea.

II Beating in stomach and bowels.

II Frequent attacks of severe vomiting, and when not so, always feeling of nausea, and fear to eat.

II Gastralgia or cardialgia, < at menses, so she would drop to the floor, comes and goes suddenly.

II Gnawing, hungry feeling, not > by eating ; everything she eats, except fish, makes her worse ; the thought of milk makes her sick.

II Nausea at beginning of diphtheria.

II Severe throbbing in region of solar plexus ; when it becomes very severe, which it did daily for hours at a time, it would seem to extend or continue upwards to head, when dizziness and lightness of head would supervene, requiring her, to lie down at once, otherwise she would fall violently to the floor. "

II Stomach tender and bloated ; cannot eat or drink anything without pain in stomach-pit.

II The almost constant diphtheritic discharges from mouth and nose nearly ceased, and she almost immediately had spells of sickness of stomach, and would occasionally vomit pieces of membrane.

I Burning in epigastric region ; feeling of weight and pressure of stone in stomach ; very thirsty ; abdomen swollen and burning, with bearing down pains therein ; mucous, yellow, liquid stools ; pulse 100 ; pains and throbbing in temples ; flushes on left cheek ; red, circular spot below right malar bone, burning to touch ; no appetite, cannot bear food ; jaw cracks while eating.

I Dyspeptic pain, as from a stone, or undigested food, in stomach-pit at 9 :45 P. m. ; followed by a stabbing pain in right lung, just below nipple.

I Nausea, with headache, on waking ; continuing all morning.

I Weak, sinking feeling at stomach-pit, on waking in morning.

Abdomen.- (Lac caninum)

II Abdomen swollen and burning, with bearing down pains therein.

II Extreme heat in abdomen.

II Feeling of tension in left groin ; does not want to walk or stand, as it < the sensation ; > by flexing leg on abdomen.

II Intense sharp pain in left side of abdomen, with nausea while leaning forward.

II Pain and burning in left side of abdomen and pelvis, with weight and dragging on that side ; clothes feel heavy.

II Pain in abdomen, < leaning forward ; > leaning back.

II Severe burning pain right hypochondriac and iliac region and corresponding part of back, extending across back to left side of abdomen ; < when on feet or when fatigued, > lying down.

II Smarting in right groin ; pains seem to be in pelvic bones, uterus and limbs.

I Abdomen very hard and swollen, in evening.

I Headache over left eye on first waking, and great pain in pelvis, most marked at right ovary.

I Pain in pelvis, principally in right ovarian region.

I Pains in abdomen intermittent.

I Pressure from within outwards, as if contents of abdomen would be forced out literally, just above pelvis.

I Very acute pain in left groin, extending up left side to crest of ilium, > by stool ; sometimes pain in track of colon.

Stool and Rectum.- (Lac caninum)

II Mucous, liquid, yellow stools.

II Obstinate constipation ; frequent desire with darting pain in rectum, no power to expel ; stool large, whitish, rough, scaly, hard.

I Constipation.

Urinary Organs.- (Lac caninum)

II Great difficulty in urinating.

II No urine for 47 3/4 hours, bladder pretty full, parts fearfully swollen, and irritation on urination very great.

II Urinating only once in twenty-four hours, and then copiously, but with some difficulty and slight irritation.

II Urine scanty, infrequent, coffee colored, no desire to urinate, quantity less than a gill in twenty-four hours, eighty per cent, albumen, with much mucus.

II Urine partially suppressed.

II Urine very scanty and dark.

I Constant desire to urinate, passing large quantities frequently ; at night she dreams of urinating, and wakes to find an immediate necessity ; a less strong and healthy person would probably have wet the bed.

I Constant desire to urinate, urine scanty.

I Constant inclination to urinate which was restrained, as urination caused intense pain when coming in contact with vulva.

I Urine dark, heavily loaded with thick reddish sediment that adhered in different colored circles to bottom and sides of vessel.

I Urine frequent ; especially at night ; scanty, high colored ; red sediment.

I Urine unusually frequent and dark.

Male Sexual Organs.- (Lac caninum)

II Buboës and chancres.

II Constant desire to urinate, with intense pain.

II Gonorrhœal pains, intermittent, in front, middle, or posterior part of urethra ; when the gonorrhœa is >, catarrh sets in.

II Great difficulty in urinating.

II Large chancre on dorsum of penis, with a fungoid bacteric mass covering whole of corona glandis, which was at first of a glossy, shining white appearance, and later covered with a fungus, looking like fully developed aphtha, edges of swollen prepuce covered with nodosities and itching.

II No urine for 47 3/4 hours, bladder pretty full, parts fearfully swollen, and irritation on urination very great.

II Small sore at entrance of urethra ; kept getting < ; prepuce involved for about half an inch, and parts of glans around urethra an open ulcer, exhaling most fetid smell, and with most excruciating pain ; hemorrhage at 10 P. M. every evening, and during day when removing dressings ; constant desire to urinate, with intense pain ; no sleep for a fortnight, red, glistening appearance.

II Urinating only once in twenty-four hours, and then copiously, but with some difficulty and slight irritation.

II Urine partially suppressed.

II Urine scanty, infrequent, coffee colored, no desire to urinate, quantity less than a gill in twenty-four hours, eighty per cent. albumen, with much mucus.

II Urine very scanty and dark.

I Chancre on prepuce, left side of frenum ; penis greatly swollen ; chancre like a cauliflower excrescence, red, smooth, and glistening, granulating rapidly from centre to circumference.

I Constant desire to urinate, urine scanty.

I Penis enormously swollen, and a chancre on glans like a cauliflower excrescence, over half an inch in diameter ; it was red, smooth and glistening ; no pain ; in a week there appeared two small chancres, deep, sharp edges, clean, and with same shining appearance.

I Prepuce involved for about an eighth of an inch, and parts of glans penis around urethra an open ulcer exhaling most fetid smell ; pain excruciating ; hemorrhages at ten every evening and during day when removing dressing ; desire to urinate constant, and accompanied by intense pain ; had not slept for a fortnight ; red, glistening appearance of ulcer.

Female Sexual Organs.- (Lac caninum)

III Great engorgement of breasts, with sensitiveness to touch, precedes menses.

III Menses ; 14 days too soon, profuse ; seven days too soon, flow came in gushes, scanty, intermittent, bright red and stringy, preceded by much flatulence from bowels ; very stringy and sticky, cannot get rid of them.

II Bearing down as though everything would fallout through vagina, with very frequent desire to urinate, and smarting in urethra.

II Constant burning pain in left ovarian region, extending from left leg even to foot.

II Foul smell from genitals.

II Great irritation about vulva and rectum.

II Great swelling of left labia and terrible pain while urinating ; from gonorrhea.

II Menses nearly ceased ; at menses much pain in right thigh and uterus, constant desire for stool, very low spirited.

II Much pain before and after menses, severe headache and entire prostration for first day or so.

II Pain in abdomen principally in right ovarian region.

II Pain in uterine region, passing down inside of thighs, half way to knees, and right leg feels numb.

II Parenchymatous metritis (two cases), in one, uterus three times as large as natural, round as a ball, and body very hard, cervix obliterated by altered form of body ; uterus sensitive.

II Pressure on labia causes a slight flow of blood ; menstruation commenced entirely normal.

II Retroverted uterus.

II Severe pain in entire uterine region, with profuse discharge of yellow, brown and bloody leucorrhoea, two weeks after menses ; intense pain and enlargement of left ovary, which could be seen protruding.

II Sharp pain in right ovary.

II Sharp, lancinating pains like knives cutting upward from os uteri, and as these were being relieved, sensation as of needles darting upwards in uterus.

II Soreness and enlargement of breasts.

II Uterine hemorrhage for six weeks ; ovarian pains alternated sides, as did the chronic headache.

I Blood bright red and stringy, hot as fire, coming in gushes and clotting easily ; constant bearing down pain, as if everything would come out of vulva.

- I Breasts seem very full.
- I Breasts very painful and sore ; feel as if full of hard lumps, very painful when going up or down stairs.
- I Breasts very painful, but no lumps ; pains are caused by least jar ; has to hold breasts firmly when going up or down stairs ; breasts < towards evening, pressure of her arm, in natural position, caused considerable pain.
- I Breasts very sore and painful, with sharp, darting pain in right ovarian region extending to knee, very painful and must keep leg flexed (1st d. after miscarriage at 6th month)
- I Breasts very sore, sensitive to least pressure ; dull, constant aching pain in them all evening.
- I Constant pain in nipples.
- I Dysmenorrhoea, abdomen sensitive even to weight of clothing ; flatus from vagina.
- I Escape of flatus from vagina.
- I In afternoon, sharp pains in right ovarian region, not constant but intermittent.
- I Inflammatory and congested condition of uterus, with extreme soreness and tenderness, that made every motion, position, and even breath painful.
- I Intense painful soreness of vulva, extending to arms, coming on very suddenly about noon, and lasting for about two hours ; came on again during evening ; could not walk, stand or sit ; > by lying on back and separating the knees as far as possible.
- I Itching in left side of labia, with rough eruptive condition on left side of vagina, with acrid leucorrhoea ; excoriating severely.
- I Itching of vulva.
- I Leucorrhoea, very profuse during day, none at night ; discharge whitish and watery ; pain in small of back ; very irritable, < standing or walking.
- I Membranous dysmenorrhoea.
- I Pain in left ovarian region ; across lower part of abdomen.
- I Pains in uterine region, all day, no particular direction except down inner side of thigh half way to knees.
- I Raw and bad-smelling sores between labia and thighs, in folds of skin ; < when walking, would rather keep still all the time ; these sores are covered with a disgusting white exudation.
- I Severe pain in right ovarian region, > by flow of bright red blood.
- I Sore throats are very apt to begin and end with menstruation.
- I Urination causes intense pain in vulva, when even least drop of urine comes in contact with it.

Respiratory Organs.- (Lac caninum)

- II Breathing hoarse and croupy, and at times an entire stoppage of breath, when it would resume with a violent effort.
- II Breathing often snoring only possible through mouth.

- || Breathing very labored.
- || Constant cough, accompanying soreness.
- || Constriction in lower part of larynx, like a finger across throat ; feeling as of a bar across back of throat.
- || Cough and dyspnea.
- || Cough caused by irritation in upper part of throat, < lying down at night, also after eating and drinking and after talking ; with soreness of left side of throat and constant desire to urinate.
- || Cough on taking a long breath, not when swallowing.
- || Croupy cough, a dry, hoarse bark, penetrating through closed doors all over the house.
- || Difficult breathing ; during evening had severe suffocating spells.
- || Excessive hoarseness, and tickling, choking sensation, moving about.
- || Expectoration of profuse, sticky, tough, white mucus in masses, with coryza.
- || Great difficulty in breathing, could not lie down flat.
- || Hard metallic cough.
- || Larynx sensitive to pressure.
- || Loss of voice.
- || Loud snoring during sleep.
- || Respiration was hoarse, crying was whispered and broken, often no sound at all.
- || Sensation as if breath would leave her when lying down and trying to sleep ; has to jump up and stir around for an hour or so every night.
- || Short breath.
- || Terrible dyspnea immediately after sleep, first on l. side of chest ; dyspnea compelled her to be lifted upright with violent exertion to get breath ; sharp pain in region of heart with each attack.
- || Throat troubles her much if she reads aloud or talks more than usual ; it seems almost as though it was stopping up, and she feels very hoarse, but has no soreness ; there is a feeling of fullness and choking.
- || Unable to speak loud ; distressed feeling while speaking.
- || Voice hoarse and husky ; interrupted by weakness and hoarseness.
- | Cough from tickling in upper anterior part of larynx, talking and lying down.
- | Tickling sensation in throat, causing cough ; in afternoon quite hoarse.

Chest.- (Lac caninum)

- || Heart beats rapidly from slight causes.
- || Pulse : quick and feeble ; 100 ; 149; 170 ; 130-140 ; almost gone.

II Pulse 117.

II Pulse 130, wiry, weak.

II Pulse 130.

II Pulse of little volume.

II Pulse rapid, quick.

II Pulse so rapid it could scarcely be counted.

II Sharp, incisive pain between scapulae, passing through sternum, with a sensation of pressure or constriction of chest in afternoon.

II Trembling, jerking and fluttering through lungs, with numb, prickling sensation all over body, legs and arms.

I Pulse quick, full and strong, with pains in chest and throat.

Neck and back.- (Lac caninum)

II Heat, pain and beating in small of back.

II Intense, unbearable pain across supersacral region, extending to right natis and down right sciatic nerve ; pain so severe as to prevent sleep or rest ; at same time diphtheritic sore throat on right side, with sensation of a lump, could not swallow solid food.

II Neck aches, making her want to bend head forwards ; entire spine sensitive.

II Neck stiff.

II Pain in back of neck.

II Pain in back.

II Spine aches from base of brain to coccyx (pharyngitis).

II Spine aches from base of brain to coccyx.

II Wandering pains in nape, with stiffness.

Upper Limbs.- (Lac caninum)

II Fingers extremely cold but not rest of hands.

II Left hand bloats and is numb, with trembling, jerking and fluttering through lungs ; numb, prickling sensation all over body, arms and legs.

II Neuralgic pains in shoulders, left then right ; then vice versa.

II Pain in left shoulder extending across to right ; could scarcely move arm.

II Pain in one or other shoulder.

II Partial paralysis of left arm, unable to raise hand to head ; on attempting to do so was seized with sharp pains in arm below shoulder.

II Sharp pain in left hand, left arm as if asleep.

II Shoulders and arms ache.

II Trembling of left hand, as in paralysis agitans.

II Two warts on little finger noticed to be leaving.

II Two warts on little finger.

I From draught in evening, sudden, violent pains in right shoulder, so much that when retiring she could not raise arm to finish toilet, as if disabled by dislocation.

I Very fetid perspiration in axilla, staining linen brown.

Lower Limbs.- (Lac caninum)

II Aching pains in limbs and back.

II After exposure to cold night air when drunk, sharp, darting pains, < by any motion, with swelling in right knee and right ankle ; next day joints of left knee and left ankle and left hand extremely painful, moderately swollen, slightly red ; ensuing day, left ankle and knee better, but right shoulder and elbow similarly affected.

II Almost constant pain in right hip.

II Articular rheumatism in right hip and knee joints, especially former ; she was seated in an arm chair, unable to move, com ; plaining of bruised, smarting, lancinating pains in both joints and in lumbar region with swelling of affected joints ; pains < by slightest motion at night ; by touch and by pressure of bed clothes ; next day pains and swelling had gone to left hip and knee joints, leaving right almost free ; the ensuing day they had almost entirely disappeared from left hip and knee joints and had again attacked right hip and knee ; complaining, moaning, and sighing on account of her sufferings and probable termination of her illness.

II Bruised pains in soles making it difficult to walk ; in twelve days pains suddenly left soles and appeared in right knee joint, being smarting, lancinating, with light swelling of joint ; could not move affected limb, as least motion < pains, as did touch and pressure of bed clothes ; on following day left knee joint affected in same way, right > ; on ensuing day right again affected, with relief to left ; afterwards hip joints attacked alternately with same symptoms, alternating like these in pains and swelling, left joints one day with > of right, and vice versa ; also lancinating pains in left side of chest ; after four days wrist joints affected, first right, with same symptoms as those of lower extremities, symptoms of one side of body alternating with those of the other ; not able to move himself in bed, lancinating pains made him cry out ; constipation, sleeplessness, no fever ; pains and swelling < every evening, night, by movement, touch and pressure of bed clothes ; numb pains chiefly in ankles <.

II Burning of hands and feet at night.

II Cannot walk any distance ; trembling through right thigh, and feeling as though entire lower portion of body was giving sway ; felt as though something was strained across lower part of bowels.

II Cramps in feet.

II Ecthyma : a sore breaks out in right leg, excessive itching ; inflammation then swelling, blisters form and suppuration sets in ; afterwards clear lymph, then discharge of matter ; then scabs and scales, turning eventually into a bran-like desquamation ; scars have left discolored skin.

II Limbs cold to knees.

II Pain in right hip and leg while walking, with a trembling of leg, and slight feeling of uncertainty, esp. on going down stairs.

II Partial paralysis of right leg from miscarriage ; has to use a cane ; leg numb and stiff, but cannot keep it still any length of time ; feels > flexing it on abdomen.

II Restlessness in legs.

II Sciatica.

I Bruised pains in soles of feet, stiffness of ankle, knee and hip joints, and occasionally intense pains which move upwards ; pain in ankle joints as of a dull plug pushing ; joints stiff and sore, tender to touch ; < from heat and least motion ; later knees and then hips became involved ; at first the left ankle was attacked, and then, after some hours suffering, the right with relief to left, and so on with knees and hips ; chest affected, terrible dyspnea coming on immediately after sleep, first on left side ; compelled her to be lifted upright with violent exertion to get breath, sharp pain in cardiac region ; urine scanty and dark ; pain in right side of throat ; generally < at 5 P. M.

I Intense, unbearable pain across supersacral region, extending to right natis and down right sciatic nerve ; pain so severe as to prevent sleep or rest (sciatica).

I Numb pains, chiefly in ankles, < while quiet, with swelling ; veins of ankles distended ; > while extreme heat is applied.

I Pains in limbs as if beaten.

I Rheumatic pains in left hip and along sciatic nerve ; wandering pains in nape of neck, with stiffness ; pains in one or other shoulder ; pain above left eye and heaviness of eyelids ; sensitiveness to light.

o

LAC VACCINUM DEFLORATUM.

Mind (Lac Vaccinum Defloratum)

I Depression of spirits ; don't care to live ; question as to quietest and most certain way of hastening one's death.

I Depression of spirits which is not dissipated by conversation.

I Head light, with throbbing in temples.

I Imagines that all her friends will die and that she must go to a convent.

Head (Lac Vaccinum Defloratum)

II General sore pain of head produced by coughing.

II Intense pain at point of exit of supra-orbital nerve, diffused thence over forehead ; attack commences with chill, quickened pulse, flushed face and belching of wind from stomach (cured).

II Throbbing frontal headache (over eyes), nausea, vomiting and obstinate constipation ; esp. in anemic women.

I After injury subject to distress in head ; severe pain in forehead just above eyes ; breath very offensive ; appetite poor, nausea ; at times sleeps for hours during attack ; great distress across back ; urine dark and thick.

I After light breakfast, pain in forehead, with nausea ; very pale face, even lips looked white ; vomiting of ingesta and afterwards of mucus and bitter water ; deathly sick feeling in pit of stomach, < rising up in bed ; profuse urination every half hour ; urine colorless as water ; great thirst ; intense throbbing pain in vertex.

I American sick headache, with gastric symptoms.

I At first a sharp pain at apex of heart, as though a knife was cutting up and down ; this lasts a few seconds and is followed by a strange feeling in head ; forehead feels extremely heavy, with dull sensation over eyes, and considerable throbbing, most marked on each side of head ; rest of head feels very light, dimness of vision ; can only distinguish light, not objects ; at same time great loss of strength ; cannot stand, but falls backwards, and remains entirely unconscious for two or three minutes ; weakness passes off gradually and is followed by crying, palpitation of heart and great depression of spirits ; imagines that all her friends will soon die and that she must go to a convent ; she can produce an attack at any time by extending arms high above head, or by pressure around waist ; spells come on at 7:30 P. M.

I Attacks come every eight days ; during attack can neither eat nor drink, nor endure light or noise ; does not even like to speak ; great prostration, < during menstruation ; when pains subside inflammation of tonsils appears ; tongue white and no relish for food.

I Dim vision, as of cloud before eyes ; profuse urination ; full feeling in head ; slight nausea at pit of stomach ; face pale ; feet cold ; coldness in back.

I Faintness and nausea when stepping upon floor in morning.

I General sore pain of head produced by coughing.

I In morning nausea and sensation of a round ball full of pain in centre of forehead (cured).

I Intense headache in forehead and through head ; < in vertex, afterward head felt bruised (cured).

I Nausea and sometimes vomiting, which > ; pain in forehead as if head would burst, with blindness ; pain is > by bandaging head tightly ; < by light and noise ; constipation, stools large ; hands and feet cold.

I Pain commencing in and above inner end of r. eyebrow before rising in morning ; soon after rising pain passed into eyeball ; < until afternoon, at which time it became unbearable ; < by walking and particularly by sitting down, though done carefully, also by heat radiated from fire or stooping, on pressure ; pressure on temples disclosed strong pulsation of artery ; pain ceased entirely at sunset and did not return till next day. Light did not < it. The eye had no unusual appearance. Lac Fel. had no effect. Partial relief by Sol. and Oxygen (cured by Lac Defloratum cm.).

I Pain first in forehead, extending through occiput, making her nearly frantic.

I Pain first in forehead, then extending to occiput, very intense, distracting and unbearable ; great photophobia, even to light of candle ; deathly sickness all over, with nausea and vomiting, < by movement or sitting up : very chilly, and external heat does not > ; frequent and profuse urination of very pale urine.

I Pains so severe that she would bury her eyes in her hands and press them into pillow.

I Periodical pain in forehead, as if head would burst, accompanied by violent effort to vomit, and more rarely vomiting ; hands and feet cold ; diarrhea alternating with constipation, the latter predominating ; loss of appetite ; smell or thought of food causes nausea ; tongue moist, coated white ; thirstlessness ; always < at menstrual period ; menses scanty and accompanied with colic.

I Severe frontal headache ; nausea and sometimes vomiting upon rising in morning, or from recumbent position at any time, or upon moving ; great constipation ; constant chilliness even when near fire ; urine profuse and watery, or scanty and high colored ; intense pain throughout whole spinal column ; excessive thirst for large quantities ; great depression of spirits ; sudden prostration of strength at 5 P. M. ; skin color of red rose, with swelling of face, neck, arms and body, generally in morning and during day and evening.

I Severe headache for years ; severe pain over eyes ; intense throbbing in temples.

I Severe headache, with a sensation as if top of head was lifted off, raised about five inches and brains were coming out ; head feels hot, motion < pain ; face felt as if flesh was off bones and their edges were separated and sticking out ; better in five minutes after taking Lac Defloratum, and next morning was entirely relieved.

I Severe pain over eyes, with intense throbbing in both temples, eyes feel as if full of little stones ; eyeballs intensely painful, and on shutting eyes, pressure of lids increases pain ; edges of lids feel contracted, and convey sensation as of a narrow band drawn tightly across eyeball ; pain over left hip ; constipation and profuse urination during paroxysm.

I Throbbing in temples.

I Vertigo : on moving head from pillow ; < lying down and esp. turning while lying, obliging to sit up.

Female Sexual Organs (Lac Vaccinum Defloratum)

II (Has never failed to bring back milk in from 12 to 24 hours.)

II °Menses suddenly suppressed by putting hands into cold water to rinse out clothes, with great pain in uterine region, intense headache, aching pains all over, fever, flushed face. 9.

II Decrease in the size of the breasts, which after the proving resumed their original size and firmness.

II Diminution of the breasts, and diminished secretion of milk in nursing women, never has failed to bring back the milk in from 12 to 24 hours.

II Menses suppressed by putting hands in cold water, with great pain in the uterine region, intense headache, aching pains all over, fever and flushed face. After 1m. in water, one spoonful at night, headache left and she slept all night. Next morning slight flow ; took another spoonful, and by 10 o'clock flow all right and she was free from pain.

Chest (Lac Vaccinum Defloratum)

II Sensible decrease in the breasts which after the proving returned to their original size and firmness.

Nerves (Lac Vaccinum Defloratum)

II Great lassitude and disinclination to exertion.

I Feels completely tired out and exhausted, whether she does anything or not ; great fatigue from walking.

I Great loss of strength, commencing with a sharp cutting pain in apex of heart ; forehead feels heavy, with a dull sensation over eyes and throbbing; principally in temples, rest of head feels light.

I Great restlessness and extreme and protracted suffering from loss of sleep at night.

Generalities (Lac Vaccinum Defloratum)

II Pain commencing across the upper sacral region, passing around either side over the hips down to groins and down thighs inside, sometimes to the feet-cured several cases.

Skin (Lac Vaccinum Defloratum)

II Bright scarlet redness on chest and around neck ; next day all over body except legs, which were, however, covered that night ; skin in large patches assumed dark red color bordering on purple, as seen in malignant cases, while body seemed swollen.

|| Crusts on skin, under which greyish yellow matter formed and was squeezed out.

|| Diphtheria with or following scarlatina.

|| Ichthyosis, with branlike desquamation of skin.

|| On face, hands, neck and chest, bright scarlet eruption exactly like scarlatina.

|| Several boils on left side.

| Herpetic eruption in both axillae, with light brownish scab, extremely painful when washing ; eruption most in r. axilla, and in both instances appeared previous to pain in labia, which was followed by a discharge of blood from vagina.

| Shining, glazed, and red appearance of ulcers on shin and wrist.

| Small blotches on chest, wrists and r. knee.

| Throat full of large foul, grey yellow patches ; deglutition especially painful after sleep and from swallowing fruits (acid) ; lumpy sensation felt in the middle of the throat ; unrest, delirium with undefined fears ; considerable bright red, fine eruption on face and chest ; itching with dry skin.

| Very small blotches like flea-bites.

Sleep (Lac Vaccinum Defloratum)

|| Aggravation of symptoms after sleep.

|| At night lies with leg flexed on thigh, and thigh on pelvis ; restless ; < after sleep.

|| Cried out and talked in sleep.

|| Dreamed a large snake was in bed.

|| Dreams of going on a journey, and was separated from party, and had to walk a long distance, and arrived at station just in time to see train start off.

|| Got to sleep late ; profuse sweat during sleep ; felt feverish all night ; in morning > in every way

|| Great desire to sleep.

|| Restless sleep at night, bad dreams.

|| Sleep prevented by being very cold for one hour after retiring, with great nervousness.

|| Sleepless and crying continually.

|| Sleeplessness from emotional strain, with entire nervous debility.

| Cannot find any comfortable position in bed ; there is no way that she can put her hands that they do not bother her ; falls asleep at last on her face.

| Dreams frequently that she is urinating, and wakes to find herself on point of doing so, requiring immediate relief.

| Sleep disturbed, very wakeful ; limbs cold all night.

I Very restless all night, could not keep clothes over her.

Fever (Lac Vaccinum Defloratum)

II Dry, hot skin.

II Fever and bathed in warm perspiration, especially about face, neck and hands.

II Fever and chills for a few days, and up and down every few hours.

II Fever returning every afternoon.

II Fever.

II High fever.

II Intense heat.

II Moderate fever.

II Perspired considerably through night, sweat having a rank smell.

II Severe chills.

II Skin clammy.

II Sweat all over.

II Temperature, 102, 102 $\frac{3}{4}$, 103, 103 $\frac{1}{4}$.

II Wakes at night in cold perspiration, with fearful foreboding.

I Exhausting sweats ; after sleep.

I Feels feverish.

I Intense fever on waking in morning, with perspiration.

Nerves (Lac Vaccinum Defloratum)

II Child partially paralyzed after diphtheria ; could not walk ; pain all over, cough, aphonia, loss of appetite, emaciation.

II Great exhaustion, with "poisoned" feeling.

II Heaviness, weakness, general languor.

II In morning so much prostrated that she could not turn in bed ; so tired.

II Often feels as if she would lose use of limbs.

II Profound depression of vitality.

II Profound prostration, to extent of refusing to make effort to take a dose of medicine.

II Sinking spells every morning, attended with great nervousness.

II Very weak.

II When walking seems to be walking on air ; when lying does not seem to touch bed.

I General weakness and prostration very marked.

I No inclination for least exertion, would like to do nothing but sleep ; much lassitude.

I Restlessness.

I Suffering from very unpleasant nervous symptoms ; not in low spirits, but weak, and nerves so thoroughly out of order that she cannot bear one finger to touch the other, and often feels as though she should lose use of her limbs ; sensation as if throat were closing, sensation is between throat and nose ; feels as if something in throat was either enlarged or relaxed, and has a desire to keep mouth open ; talking difficult ; disposition to talk through nose ; sometimes cannot swallow, because there seems to be a kind of muscular contraction in throat ; sleep restless, frequently wakes with sick headache, which seems to commence at nape ; wakes with severe pain at lower part of back ; pain leaves when about work a short time, does not return until next morning ; nerves very much overwrought, afraid of being unable to perform duties.

○

LAC VACCINUM (Cow's Milk).

Stool (Lac Vaccinum)

II Obstinate constipation ; stool hard, dry ; in impacted balls ; can be passed only with great straining.

○

LYSSIN (Hydrophobinum ; saliva of a rabid dog).

Mind (Lyssin)

II Lyssophobia ; fear of becoming mad.

II When he hears water poured out, or if he hears it run, or if he sees it, he becomes very irritable, nervous ; it causes desire for stool and other ailments.

I A musician received a bite from a small pet dog in calf of left leg while walking through a dark entry ; the bite was very slight, scarcely wounding skin ; the animal was in a healthy condition and remained so ; a pain in bitten place kept returning from time to time independently of mind dwelling on it, until it finally grew to a burning which extended through entire body, causing an indescribably strange sensation ; in night trembling, and a tormenting fear that he would have hydrophobia ; was thirsty and drank water freely ; it was two years after the bite when these and the following symptoms appeared ; frequent spitting of saliva all through day, only ceasing awhile after taking strong tea in evening ; disturbed dreams at night ; finally could not partake of food or drink, and complained of pricking stitches under tongue ; mental excitement now as all his life, affects him badly ; hot vapor relieved burning pain in bite, and Lyssin 2C. (Jenichen) improved all symptoms rapidly ; in three weeks he considered himself cured, and started on a journey.

I Attacks of nervous headache become awful and insupportable if he hears water run out of a hydrant.

I Break out of their stables furiously and run and jump over ditches and fences. Lyssa of sheep.

I Exhilarated, felt as if he had received joyful intelligence.

I Feels as if he cannot physically endure his fears much longer, and shall be compelled to go into an insane asylum.

I Felt same rheumatic pain his brother complained of.

I Frequent gaping without being sleepy, most when he has to listen to others.

I Ill humor.

I Inclination to be rude and abusive, to bite and strike.

I Mania spermatica ; stallions.

I Mary M., aet. 17, had been bitten several years previously by a dog, and reading of several cases of hydrophobia as reported in papers, was found in following condition ; crouched in corner of sofa, dark red bloated face, expression of terror in face, eyes glistening, conjunctiva injected red ; was brought into this condition by endeavoring to take a drink of water, and could not hear water mentioned without a shudder of fear, could not swallow, pulse very high, tongue dry and coated red ; Lyssin 200 one dose ; she was better next day, but had several slight attacks afterwards, always induced by running of water, but always yielding to remedy ; has not had an attack in nearly a year.

I Memory for single words much improved.

I Mental emotion always makes him worse.

I Not afraid of dogs, but dislikes to see them because their sight renews her fear. (Lyssophobia, after bite by non-rabid dog.)

I Restless, constant bleating in a hoarse voice.

I Sometimes he could control inclination to stool by a strong effort of will, but effort caused much nervous irritation.

I Strange notions and apprehensions during pregnancy.

Head (Lyssin)

II Beating, throbbing headache ; most severe in r. temple and above r. eye ; each bone feels shattered and sore ; from temple to temple.

II Burning aching from left side of occiput down neck.

II Dizziness : as if something was drawing around in a circle, and as if she could not hold her head straight ; after lying down, like a shock in upper part of brain. ; with inclination to fall to r. when stooping ; towards evening in upper part of head, as if she would fall, while walking ; frequent and transient ; with dim sight while walking and sitting ; and nausea ; with cramps in abdomen ; temporarily > by return of diarrhea ; < by teaching, spelling, or being obliged to notice letters ; after lying down in bed, shock in upper part of brain ; dimness of vision on rising from stooping ; on rising from chair, staggers ; while sitting ; on rising, cannot walk straight.

II Frequent pressure on vertex, as if a cast, which fitted top of head, was pressing it down.

II Headaches from bite of dogs, rabid or not.

II Peculiar lightness in head ; lightness after nausea.

I Afternoon, pain in head ; sick headache.

I At noon slight headache, lasting all day.

I Maddening outward pressing pain in forehead ; he presses his head against the wall.

I Pain as if head would burst.

I Rush of blood to head : while lying down ; from chest upward, with toothache ; during pregnancy ; when rising.

I Violent headache and backache.

I Violent headache, most in temples and forehead, < during day and from stooping and stirring about.

Eyes and Sight (Lyssin)

II Swelling of eyelids after bite of dogs. In sheep.

I Could not see or hear.

I Extreme ulceration of eye, lids closed and puffed up by pus. In sheep.

I Great inflammation in an eye from which gushes foamy pus ; pustules around eye ; in morning lid puffed up like an eggshell, small pustules on one finger.

I Inflamed, dim, watery, staring eyes, with very much dilated pupils, upper lid drawn up, and diminished sight.

I Sight of water : agitation ; renews idea of pain ; causes convulsions (pregnancy).

I Since five years could not read longer than a few minutes at a time, when she would see letters double and would read something else than the right words ; had tried all sorts of spectacles without benefit, was often ashamed because she could not sign her name properly.

I Sore eyes and some fever after a bite in nose.

Ears and Hearing (Lyssin)

II Hearing water poured out in next room makes him very irritable and nervous.

I After the pain following an evacuation had lessened, and he was sitting at open window, the large street water plug was opened to cleanse the streets, and as soon as he noticed water running down gutter in front of his house, he was seized with violent pains and had to go at once to the water closet.

I During his attacks of chronic headache, which come after mental emotion or excessive mental exertion and last a day or two, he cannot bear to hear running water ; if hydrant is allowed to run in an adjoining room, or even if water is poured into a basin, his headache increases to an insupportable degree.

I Hearing water poured out brought on convulsions.

I If, during night, or in morning before rising, he heard pouring out of water in next room, he was immediately obliged to rise and have an evacuation.

Face (Lyssin)

II Masseter muscles not affected by spasms.

I Appearance peculiar ; skin sallow, pale or anemic ; bloated all over, but no "pitting."

I Pain in left cheek near nose, from fifth to seventh day ; followed by a hard, painless swelling of size of a pea, turned redder on thirtieth day, and became covered with a small scab ; on thirty-second day it grew soft and turned bluish red, particularly around edges ; on puncturing, a disagreeable yet not severe pain at some distance from swelling and more inside of cheek ; a small quantity of blood and matter was discharged ; on fortieth day the place ceased to discharge and healed.

Teeth and Gums (Lyssin)

I Aching in swollen gums on r. side.

I Grinding of teeth.

I Toothache and other complaints during pregnancy, with internal ebullition of blood from chest to head ; head feels as if filled with air to bursting.

Taste and Tongue (Lyssin)

II Difficult, incorrect speech.

I Impediment in his speech, would begin a sentence with difficulty after several fruitless attempts ; some palatal vowels he could not pronounce, others but incorrectly.

I Pricking sensation under tongue.

I Tongue large, pale and flabby.

Mouth (Lyssin)

II Frequent spitting.

I At 10 A. M. began to spit a great deal, continuing all day till supper.

I Constant sensation of intense dryness of mouth and throat.

I Large quantities of tough saliva in mouth, with constant spitting.

I Spits all the time small quantities of a frothy saliva ; with pain in limbs.

I Tough, short frothy phlegm in mouth (horse).

Throat (Lyssin)

II Sore throat, constant desire to swallow ; much saliva and feeling as if beaten.

I A slight redness of affected parts.

I Burning stinging in throat.

I Constant desire to swallow, painful and ineffectual.

I Difficulty in swallowing ; particularly fluids.

I Had difficulty in speaking ; a connected sentence could only be uttered after making several fruitless attempts ; palatic letters could not be pronounced at all, all were pronounced wrong.

I Periodical spasm of esophagus, continual painful inclination to swallow without being able to swallow anything ; constriction is most severe when taking water into mouth, if he tried to swallow it forcibly, he had burning and stinging pain in throat, cough and retching which forced fluid from his mouth ; difficult speech.

I Periodical spasms of esophagus.

I Slight redness of palate and throat, with spasm of esophagus and difficult speech.

I Sore throat, as after swallowing red pepper.

I Trying to swallow water forcibly, it caused burning and stinging in throat, besides cough and gagging, which forced out contents of mouth.

I When taking water into mouth, constriction was greatest.

Appetite, Thirst, Desires, Aversions (Lyssin)

I Abnormal cravings during pregnancy.

I Appetite good, but digestion deficient ; a portion of nearly everything he ate passing bowels in an undigested state.

I Aversion to drinking water, but can take small quantities of chocolate.

I Aversion to water ; imagines he cannot swallow ; says he is thirsty, but cannot look at water or hear it poured out.

I Could take no nourishment (with spitting), could not remain at table.

I Drank some strong tea at supper, after which salivation discontinued until 8 P. M., then commenced again.

I Excessive desire for salt.

I Felt very thirsty and had no aversion to drink, on contrary drank large quantities of water.

I Had not been drinking any water for some time, only some hot tea.

I Want of appetite.

Hiccough, Belching, Nausea, and Vomiting (Lyssin)

I Gagging when he forcibly attempts to swallow water, forces it out of his mouth.

Hypochondria (Lyssin)

I Painful throbbing as if an abscess was forming in region of spleen, but very deep in, exact locality is half way between median line and outline of left side ; it lasted eight days ; with it departed remnant of a similar affection in this locality, against which eleven years of allopathic treatment had proved of no avail.

Abdomen (Lyssin)

I General soreness in whole of lower abdomen.

Stool and Rectum (Lyssin)

II Dysenteric stools with tenesmus ; renewed as soon as he hears or sees water run.

II Stools of bloody mucus.

I Chronic diarrhea contracted in Southern camps.

I Stools watery and profuse, with severe pains in lower bowels ; frequency of stools not uniform, some days five or six, others fifteen or twenty, usually more frequent in morning.

I Straining to evacuate, causing a violent pain in small of back and in rectum, afterwards compelling him to walk about, although weak.

I Tenesmus during and after stool.

I When he sees or hears running water, violent pain and tenesmus return ; in a case of dysentery in Summer, of six weeks standing, in which stools were most frequent in night, consisted of bloody mucus and were followed by pains in rectum and small of back, which forced patient to walk about in spite of great weakness ; could neither stand nor lie down ; had drank no water whole time, only hot tea occasionally.

I When in morning some water was poured out from pitcher into basin, pain and desire to stool returned.

Urinary Organs (Lyssin)

I Constant desire to urinate on seeing running water ; urinates a little at a time.

I Urine too scanty and high colored.

Male Sexual Organs (Lyssin)

II Complaints resulting from abnormal sexual desire.

I Atrophy of testicles ; testicles diminish in size, first left, then right.

I Hydrocele.

I Increased sexual desire. (Dropsy of spine with sheep. Hydrophobia of sheep.)

I Satyriasis in a stallion ; hot breath streamed from nostrils.

I Semen is discharged too late or not at all during coition.

I Strong erections, without sexual excitement or thoughts, in evening while undressing in cold room.

Female Sexual Organs (Lyssin)

II Sensitiveness of vagina rendering coition quite painful.

I Abrasion about os tincae (treated with caustic) some tumefaction of cervix and walls of vagina remained, showing a low degree of inflammation.

I After menses found that prolapsus uterus, a case considered incurable, was in its proper position, and continued so after an interval of two months, though she has done much to test the cure, lifting a heavy child in and out of bed at night, when she was necessarily without her supporter.

I Anemia in consequence of disturbed sexual function.

I Bright redness of vaginal portion of womb.

I Considerable pain in lower part of back, with a soreness felt through pubic region, which was clearly proved to be in neck of womb by an increase of pain from pressing finger on neck ; principally at point where finger came in contact with womb.

I During interval of menses "a show" that seemed quite obstinate.

I Insatiable heat ; with cows.

I Menses rather frequent, protracted, dark, and at times fetid.

I Menses too profuse, at times a little too frequent.

I Menstruation, with hemorrhoids, pulsations in anus and weakness of back.

I Metritis, prolapsus or induration of uterus ; in cows.

I Prolapsus uteri of seven years' standing.

I Severe leucorrhoea, with pains in back and lower part of bowels ; sore vagina.

I Speculum showed os tincae of size of a small goose quill, smooth and normal, except that there was a string of bloody mucus of size of os hanging from it, so tough and viscid that it was difficult to wipe it away with a sponge.

I Swelling of womb in all its parts, extending somewhat into vagina. Prolapsus.

I Tearing, followed by pressing downward, could not make a hard step during catamenia.

I Weakness in back, with copious catamenia.

I With a painful sensitiveness of womb, slight degree of prolapsus, so that after any considerable effort there would be a strong conviction that it was prolapsed.

I Womb high up in abdomen, enlarged in fundus.

Pregnancy. Parturition. Lactation (Lyssin)

II During pregnancy : strange notions, desires or cravings ; rush of blood from chest upward ; toothache, backache and other complaints ; great sense of bearing down ; intense pain from inflammation of os and cervix (formerly treated with Caustic.) ; great soreness in lower part of back and bowels.

I All changes of position that tilt or rotate to a moderate extent the os uteri cause much pain.

I Since cessation of lochia a severe leucorrhoea ; pain in back and lower part of bowels ; soreness of vagina.

I Since childbirth more pain with coition and a dislike to it.

I Spasms excited whenever she attempts to drink water, or if she hears it poured from one vessel into another ; sight or sound of water affects unpleasantly, even though desiring water.

Voice and larynx. Trachea and bronchia (Lyssin)

I Pain next to larynx, on r. side, felt on turning neck and on pressure.

Respiration (Lyssin)

II Sighing with pain in heart.

I Dyspnea : with flatulency, cough and rattling in chest ; with sighing, groaning respiration ; from cardiac pain ; < lying down.

Cough (Lyssin)

I Cough and gagging.

I Coughing when attempting to swallow water forcibly, forces it out of mouth.

Inner Chest and Lungs (Lyssin)

I Gangrene of lungs, caused by pneumonia and sexual excitement.

Heart, Pulse and Circulation (Lyssin)

I Heart had for three months not been free from a sticking, drawing, squeezing pain, result of an attack of rheumatism and cold, together with a palpitation and difficulty of breathing.

I Pain in cardiac region, to which he is subject, is < half an hour after, but much > in several days.

I Stitches in heart, more while walking ; they would kill him if they continued.

Neck and Back (Lyssin)

I A sore lameness in back, with some degree of soreness in lower abdomen.

I Backache and headache.

I Considerable pain in lower part of back, with a soreness felt through the pubic region ; pressing finger on neck increased it.

I Disease peculiar to sheep, termed the "gid ; " dropsy of spine.

I Great weakness in back, as if it would split and fall apart.

I Sheep turn and wriggle their backs because they cannot bear heat of sun ; increased sexual excitement ; symptoms pointing to madness ; scratching their backs appears to be agreeable to them, as they remain quiet and make a peculiar motion with their mouths, which does not express pain.

Lower Limbs (Lyssin)

II A pressive pain in r. hip bone, goes from there to middle of sacral bone.

II Ulcers on legs, with emaciation and coldness, left side ; contraction of hamstrings and relaxation of ligaments of ankle joint and flexors of toes.

I Great swelling in heel of a cow from dog bite ; very restless.

I The pain in calf came even when he was not thinking of bite at all ; lately much < and quite severe.

Nerves (Lyssin)

II Attacks returned every few minutes, for five hours, until they were arrested ; pain passing down spine to loins and hips, and from thence to knees.

II Convulsions daily ; rolls head from side to side ; winking and rolling eyes ; attempts to bite others ; pelvis and legs are turned to one side as far as possible. Child, aet. 2, after scarlatina.

I At times a singular sensation, a kind of "die away" feeling, quite instantaneous, cannot describe it ; seems to extend through heart to back ; accompanied with quivering around breast bones and pain in nerves, from left side of chest to throat and left jaw ; fluttering about heart.

I Continual trembling sensation through whole body.

I He could neither lie down nor stand up for any length of time.

I The strange feeling changed during night into a trembling, and he is full of fear.

If during great restlessness she attempts to sit down and work, she has alternately twitching in arms and legs.

Sleep (Lyssin)

I During night slept but little ; was disturbed by disagreeable, disconnected dreams ; felt strangely. Lyssophobia.

Temperature and Weather. (Lyssin)

I Burning sensation in bitten place > by hot steam striking against it. Lyssophobia.

I Unbearableness of heat of sun.

Fever (Lyssin)

I Intermittent fever.

Attacks, Periodicity (Lyssin)

I First day > ; next day < ; on third day salivation returned. Lyssophobia.

I Periodically since two weeks.

Tissues (Lyssin)

I Chlorosis.

Touch. Passive motion. Injuries (Lyssin)

I A bite of an angry dog into r. thigh had not healed, turned into malignant ulcers ; surrounding bluish red edges, raised and hard, bases badly suppurating and ichorous, with redness and hardness.

I A cow bitten by a dog in foot had a large swelling of part, and animal got very restless.

I Felt a continued pain in spot on r. calf where he was bitten, but without a lesion. Lyssophobia.

I Pain on spot bitten became burning, and extended over whole body, and he felt very strangely. Lyssophobia.

I Scars from a bite in nose swollen and red for fifteen months.

I Two puppies, bitten by their rabid mother, did not get mad after taking Lyssin.

I Ulcers remaining after bite of evil disposed dogs.

I Wounds from bite of dogs, in sheep.

Skin (Lyssin)

I Bluish discoloration of bitten place (after Laches.).

I Cancerous sores.

I Malignant ulcers from bite of a dog.

I Pustules : on forehead ; around inflamed eye ; on finger (after bite).

I Red scar from bite of a dog.

- o **MAGNETIS POLI AMBO (The Magnet).**

Mind and Disposition (Magnetis Poli Ambo)

II Hurried heedlessness and forgetfulness ; he says and does something different from what he intends, omitting letters, syllables, and words.

II While attending to his business in the daytime, he talks aloud to himself, without being aware of it.

I He endeavors to do things, and actually does things contrary to his own intentions.

I He inclines to be angry and vehement ; and after he has become angry, his head aches as if sore.

I Hurried zeal ; afterwards a gnawing pain in the arm and in the head of the humerus.

Head (Magnetis Poli Ambo)

II Headache occasioned by the least chagrin, as if a sharp pressure were made on a small spot in the brain.

I Pain in the region of the vertex, at a small spot in the brain, as if a blunt nail were pressed into the brain ; the spot feels sore to the touch.

Face (Magnetis Poli Ambo)

II Sweat in the face without heat, early in the morning.

Eyes (Magnetis Poli Ambo)

I Dilated pupils with cheerfulness of the mind and body.

I Fiery sparks before the eyes, like shooting stars.

I Itching of the eyelids and eyeballs in the inner canthus.

Teeth and Jaws (Magnetis Poli Ambo)

I Uniform pain in the roots of the lower incisors, as if the teeth were bruised, sore or corroded.

Mouth and Pharynx (Magnetis Poli Ambo)

II Ptyalism every evening with swollen lips.

I Bad smell from the mouth which he does not perceive himself, also with much mucus in the throat.

I Continual fetid odor from the mouth, without himself perceiving it, as in incipient mercurial ptyalism.

Taste And Appetite (Magnetis Poli Ambo)

II Hunger, especially in the evening.

I Aversion to tobacco, although he relished it.

I He has an appetite, but the food has no taste.

Abdomen (Magnetis Poli Ambo)

II Loud rumbling in the abdomen, early in the morning when in bed ; afterwards colic as if from incarceration of flatulence.

II Qualmish sensation and painfulness in the intestines, as if one had taken a resinous cathartic or rhubarb, with painful emission of hot, putrid flatulence.

Stool (Magnetis Poli Ambo)

II Blind hemorrhoids after soft stool, as if the varices on the margin of the anus felt sore, both when sitting and walking.

II Constipation as if the rectum were constricted and contracted.

II Itching hemorrhoids.

II Prolapsus recti when going to stool.

II Violent hemorrhoidal pain in the anus after stool, erosive as if sore, and as if the rectum were constricted.

I Burning at the anus when sitting, as in hemorrhoids.

Male Genital Organs (Magnetis Poli Ambo)

II Early in the morning he feels a burning in the region of the vesiculae seminales.

II Menses had ceased a few days before, returned next day after imposing the magnetic surface and continued ten days.

II Nightly emissions of semen.

II The prepuce retreats entirely behind the glans.

II Want of sexual desire, aversion to an embrace.

I The penis remains in a relaxed condition, in spite of all sexual excitement.

I Violent continuous erections, early in the morning when in bed, without any sexual desire.

Larynx (Magnetis Poli Ambo)

II Mucus in the trachea which is easily hawked up, evening and morning.

II Violent fit of cough, with profuse expectoration of blood.

I Convulsive cough.

Chest And Lungs (Magnetis Poli Ambo)

I Spasmodic cough, with shocks in the chest and anxious breathing, and visible oppression of the chest.

Back (Magnetis Poli Ambo)

II Pain in sacro-lumbar articulation, in the morning when in bed lying on the side, and in daytime when stooping a long time.

I Shock or jerk in the small of the back, almost arresting the breathing.

Upper Limbs (Magnetis Poli Ambo)

II Drawing from the head down to the tip of the fingers.

I Deep-seated pain in the arm, extending as far as the elbow, the arm going to sleep and trembling spasmodically.

I Prickings in the arm.

Sleep (Magnetis Poli Ambo)

II Early in the morning he sleeps on his back, one of his hands lying under the occiput, the other over the region of the stomach, the knees being apart, with moaning inspirations, half-opened mouth, and low muttering, dreaming of amorous things and emissions, although no emission takes place ; headache in the occiput after waking, as after an involuntary emission of semen, with asthma, and bruised pain in all the joints, going off after rising and during motion, with copious expulsion of catarrhal mucus.

II He wakes at three o'clock in the night ; in a few hours he falls into a sopor full of dreams ; afterwards, feeling of heat in the limbs without thirst, the limbs requiring to be uncovered at first, but afterwards covered carefully.

I Dreams full of feasting, boasting, and bragging.

I Lascivious dreams, even during the siesta, with discharge of the prostatic fluid.

I Vivid dreams, as if he saw the thing taking place in his waking state.

General Symptoms (Magnetis Poli Ambo)

II Shuddering movement through whole body.

II The trunk, while in a recumbent position, is jerked up spasmodically as it were, the head being jerked forward upon the bed, and then again backward upon the cushion.

I Dull, numb pain.

I Jerking shock, causing the trunk to bend violently upward and forward as low down as the hips, with cries.

I The joints are painful when touched.

I The place where the magnet had been imposed goes to sleep, becomes numb and insensible.

I The upper part of the body is spasmodically lifted and jerked forward, and then again backward on one side.

•

MAGNETIS POLUS ARCTICUS (North-pole of the magnet).

Mind And Disposition (Magnetis Polus Arcticus)

II Calm, composed mood, devoid of care.

II He makes mistakes easily in writing.

II Indolent mind.

II While attending to his business, he talks aloud to himself.

I Hasty, bold, quick, firm.

I Indolent fancy ; he sometimes felt as if he had no fancy at all.

I Sadness, in the evening ; he had to weep, contrary to his will, after which his eyes felt sore.

Head (Magnetis Polus Arcticus)

II Pushing tearing in the head behind the left ear when sitting.

II Rush of blood to the head, and suffusion of heat in the cheeks.

II Sensation as if the head were pressed down by a load.

I Aching pain over the left temporal region, externally.

I Disagreeable, compressive sensation in the head, and as if one part of the brain were pressed in.

I Drawing-boring pain in the right temple, accompanied with a spasmodic pain below the right malar bone.

I Headache, as if the temples were pressed asunder.

I Violent headache the whole afternoon, as if the brain were pressed asunder.

Eyes (Magnetis Polus Arcticus)

II Burning in the weak right eye ; it became red and filled with water (the magnet being held in contact with the weak right eye for a quarter of an hour).

II Coldness in the weak eye for three or four minutes (the magnet being held in contact with that eye for two minutes).

II Drawing in the eyelids with lachrymation.

II Excessive lachrymation ; the light of the sun is intolerable.

II Itching in the inner canthus and in the margin of the eyelids.

II Jerking and drawing in the eyelids.

II Painful feeling of dryness in the eyelids in the morning on waking.

II Pricking in the eyelids.

II Sensation as of a cobweb in front of the eyes.

II Staring look.

II Strong drawing over the eye, in the surface of the cheek, ear, extending into the upper maxillary bone (the magnet being in contact with the eye).

I Formication between the two eyes.

I Lachrymation early in the morning.

I The eyes protrude.

I Uneasy motion of the eye, with a good deal of water accumulating in either eye.

Ears (Magnetis Polus Arcticus)

II Whizzing and a drawing sensation in the ear.

I A kind of deafness, as if a pellicle had been drawn over the right ear, after which heat is felt in the ear.

I Fine ringing in the opposite ear (immediately).

I Ringing in the ear of the same side.

Nose (Magnetis Polus Arcticus)

II Illusion of smell : he imagined the room smelled of fresh whitewash and dust, he imagines the room smells of rotten eggs, or of the contents of a privy.

II Redness and heat of the tip of the nose, followed by hot, red, circumscribed spots on the cheeks.

II Violent bleeding at the nose, for three afternoons in succession, increasing every afternoon, and preceded by an aching pain in the forehead.

Jaws And Teeth (Magnetis Polus Arcticus)

II The toothache ceases when walking in the open air, and returns in the room.

II Throbbing in the tooth, with burning in the gums, and swollen, red, hot cheeks, with burning pain and heating in the cheeks, in the afternoon.

II Toothache in the direction of the eye, a very quick succession of pickings in the hollow tooth, with swollen inflamed gums and a red and burning cheek ; the toothache increased very much immediately after a meal, improved when walking in the open air, but aggravated in a smoky room.

I Drawing-aching pain coming from the temple, below the mastoid process, between the sterno-cleido-mastoideus muscle and the ramus of the lower jaw.

I The toothache is worse after eating and in the warm room.

I Throbbing in the hollow tooth (immediately), followed by a pressure in the tooth as if something had got into the tooth, with drawing in the temples.

I Toothache as if the tooth would be torn out, worse after a meal, and when sitting or lying down, improving when walking.

Mouth (Magnetis Polus Arcticus)

I Painful humming in the hollow teeth of the lower jaw, worse on the right side, the toothache ceases during eating.

Stomach (Magnetis Polus Arcticus)

II Frequent eructations of mere air.

Abdomen (Magnetis Polus Arcticus)

II Gurgling in the abdomen as if a quantity of flatulence were incarcerated, causing a writhing sensation, which rises up to the pit of the stomach, and causes eructations.

II Inguinal hernia.

I Boring pain above the left abdominal ring, from within outward, as if hernia would protrude.

I Flatulent colic immediately after supper ; sharp pressure in every part of the abdomen from within outward, as if the abdomen would burst ; relieved when sitting perfectly still.

I Relaxed condition of the abdominal ring, increasing from day to day ; hernia threatens to protrude, especially when coughing.

I She is waked in the night by a pressure in the abdomen as from a stone.

I Sore pain in the abdominal ring, when walking.

I Spasmodic contractive sensation in the hypogastrium, externally and internally, early in the morning.

Stool (Magnetis Polus Arcticus)

I Drawing, almost dysenteric pain in the hypogastrium, early in the morning, followed by difficult expulsion of the very thick feces.

I Stinging-pinching in the rectum.

Genital Organs (Magnetis Polus Arcticus)

II Nightly involuntary emission.

Chest (Magnetis Polus Arcticus)

II Pressure in the region of the heart (immediately).

I Sudden oppression of the chest, with anxiety.

Back (Magnetis Polus Arcticus)

I Gurgling and creeping sensation between the scapulae.

I Pain as if bruised in the left shoulder-joint both during motion and rest, painless when touching it.

Upper Extremities (Magnetis Polus Arcticus)

I Cramp-like sensation in the arm, and as if it had gone to sleep.

I Painful and almost burning itching in the dorsum of the middle phalanx of the little finger, as if the part had been frozen ; the place was painful to the touch.

I Prickling pain in the arm as far as the shoulder, especially in the long bones of the forearm.

I Sore pain in the right shoulder when walking in the open air.

I Violent coldness in the arm over which the magnet had been moved (in a female in magnetic sleep, after being touched with the north pole magnet).

Lower Extremities (Magnetis Polus Arcticus)

I Burning pulsative stitches in the calf.

I Excessive weakness of the lower limbs, when walking, as if they would break.

I Painful crawling in the toes of the right foot.

I Rigid tension in the hamstrings, when rising from a seat, as if too short.

I Sore pressure in the corns, which had been painless heretofore, when pressing the feet ever so little.

I Sudden lancinations in the heels, big toe and calf when sitting.

I Tearing with pressure in the outer side of the knee down to the outer ankle.

I Voluptuous itching of the left thigh, on the anterior and internal surface, but more on the former.

I Voluptuous itching under the toes of the left foot.

Sleep (Magnetis Polus Arcticus)

II Restless sleep ; he tosses about and his bed feels too warm.

I At two o'clock in the morning he is half awake, his internal consciousness being very intense, his memory very vivid and his mind filled with ideas ; he composes in the very best style in a foreign language which he neither spoke nor wrote fluently, as if he had been in a magnetic sleep ; but is unable, when waking, to recollect distinctly what he had thought.

I Constant drowsiness in the day-time.

I Excessive spasmodic yawning with pain in the articulation of the left jaw, as if it were on the point of being dislocated.

I Lascivious dreams the whole night.

I She saw a person in a dream, and next day she saw that person in reality for the first time.

Fever (Magnetis Polus Arcticus)

I Cool sweat all over.

I Heat in one of the cheeks, accompanied with a feeling of internal heat, irritable disposition and talkativeness. Sensation of warmth in the feet.

I Sensation of coldness or coolness over the whole body, as if she were dressed too lightly, or as if she had taken cold, without shuddering ; immediately after she had a small loose stool which was succeeded by pressing.

I Shuddering all over at the moment when the north pole was touched by the tip of the tongue.

Skin (Magnetis Polus Arcticus)

I Crawling over the skin.

General Symptoms (Magnetis Polus Arcticus)

II Sensation of coldness in the part which was touched by the magnet.

II Tremor in the part touched by the magnet (immediately).

II Tremulousness through the whole body, especially in the feet.

I Bruised pain in the adjoining parts, and as if one had carried a heavy burden.

I Continuous digging-up stitches in various parts, becoming sharper and more painful, in proportion as they penetrate more deeply into the flesh.

I Tensive sensation in the adjoining parts.

•

MAGNETIS POLUS AUSTRALIS (South Pole of the Magnet).

Sensorium (Magnetis Polus Australis)

I Vertigo as if intoxicated, as if he were obliged to stagger, some vertigo even while sitting.

Head (Magnetis Polus Australis)

I Drawing-tearing pain in the left brain, resembling a slow, burning stitch.

I Heaviness of the head, with a sort of creeping or fine digging in the head.

I Pressure in the occiput, in alternate places.

I Rush of blood to the head, without heat.

Eyes (Magnetis Polus Australis)

II Painful, smarting dryness of the eyelids, especially perceptible when moving them, mostly in the evening and morning.

II Watery eyes from time to time.

I Deficient sight : things looked dim, also double, when touching the nape of the neck.

Ears (Magnetis Polus Australis)

I Inflammation of the outer ear, the grooves of that portion of the ear assuming the appearance of sore rhagades.

I Noise in the ears, like the motion of a wing.

I Occasional stitches and ringing in the ear.

I Roaring in the ears, which he felt more in the upper part of the head.

I Sensation as of the whizzing of the wind in the ears, early in the morning ; he feels it as far as the forehead.

I Tearing pains in the cartilages of the outer and inner ear, extending very nearly as far as the inner cavities.

Jaws And Teeth (Magnetis Polus Australis)

I Tearing jerking in the upper jaw towards the eye, in the evening.

Throat (Magnetis Polus Australis)

I Burning in the pharynx, a sort of strangulation from below upwards, with a feeling of heat.

Appetite and Taste (Magnetis Polus Australis)

I Indifference to milk, bordering on aversion, early in the morning.

Abdomen (Magnetis Polus Australis)

II Emission of a quantity of flatulence.

I Distended abdomen in the evening immediately before going to bed, with colicky pains.

I Drawing pain in the right side of the abdomen, scarcely permitting him to walk.

I Flatulent colic at night : portions of flatulence seem to spring from one place to another, which is painful, and causes a disagreeable grumbling sensation, or a sore pinching pressure from within outward in many places, depriving him of sleep ; short flatus goes off now and then with pain, but affords no relief.

I Loud rumbling in the abdomen.

I Tearing colic occasioned by (reading ?) and walking, and appeased by sitting, especially in the epigastrium (early in the morning).

Urinary Organs (Magnetis Polus Australis)

II Incontinence of urine.

I Smarting pain in the forepart of the urethra, during the emission of urine, as if the urine were acrid or sour.

Male Sexual Organs (Magnetis Polus Australis)

I Drawing in the spermatic cord, early in the morning when the testicle is hanging down, as if pulled or distended ; the testicle is even painful to the touch.

I Impotence : embrace with the proper sensations and erection ; but at the moment when the semen is about to be emitted, the voluptuous sensation is suddenly arrested, the semen is not emitted, and the penis becomes relaxed.

I Jerking in the spermatic cord.

I Nocturnal emission (in a person affected with hemiplegia) ; it had not taken place for years past. (Note by Hahnemann. - After this emission the paralysis became worse ; the sick limb seemed dead to him.)

I Red spot, like a pimple, on the corona glandis and on the internal surface of the prepuce, without sensation.

I Slow, fine, painful drawing in the spermatic cord.

I Spasmodic drawing up of the testicles, in the night.

I The glans is red and inflamed, with itching and tension.

Female Sexual Organs (Magnetis Polus Australis)

I The menses, which had already lasted the usual time, continue to flow for six days longer, only during motion, not when at rest ; every discharge of blood is accompanied with a cutting pain in the abdomen. (Note by Hahnemann.-This woman held the south pole, touching at the same time the middle of the bar. The south pole appears to excite hemorrhage, and especially from the uterus, as its primary effect ; the north pole seems to act in the contrary manner.)

Respiratory Organs (Magnetis Polus Australis)

I Shortness of breath in the pit of the stomach.

Chest (Magnetis Polus Australis)

II Palpitation of the heart.

Back (Magnetis Polus Australis)

I Gnawing and smarting in the back.

Upper Extremities (Magnetis Polus Australis)

II Quick, painful jerking, in the arms, from above downward.

I Beating in the finger in contact with the magnet.

I Crawling in the left arm, from above downward, resembling small snakes.

I Jerking in the fingers which are touched by the magnet.

I Sense of heat and jerking in the finger touching the magnet.

I Sense, as of rumbling and gurgling, down the left arm (Immediately).

Lower Extremities (Magnetis Polus Australis)

III Soreness of the inner side of the nail of the big toe in the flesh, as if the nail had grown into the flesh on one side ; very painful, even when slightly touched.

II Pinching occasioned by the shoes on top and on the sides of the toes, and near the nail of the big toe when walking, as from corns.

I Cracking of the knee-joint during motion.

- I Drawing pain in the outer side of the bend of the knee.
- I Drawing, with pressure, in the muscles of the thighs, worse during motion.
- I Itching-burning, slow stitch in the side of the calf.
- I Sense of coldness in the right thigh.
- I Tearing with pressure in the patella (worse during motion), and aggravated by feeling the part.

Sleep (Magnetis Polus Australis)

- II Dreams about fires.
- I He quarrels and fights in a dream.
- I Restless, frequently turns from side to side, in the night when in bed.
- I Sleepless and wakeful before midnight, and no disposition to go to sleep.
- I Unusual beating in the region of the heart.

Fever (Magnetis Polus Australis)

II During the chilliness, or the feeling of coldness, he was quite warm, but he was obliged to lie down, and to cover himself well ; his mouth was very dry ; afterwards he was covered with a profuse sweat all over, without feeling hot ; on the contrary, he felt a constant shuddering over the perspiring parts, as if they were covered with goose-skin ; accompanied with a sensation as of a breeze blowing into the ears.

- I Chilliness of the legs up to the knee, with ascension of heat and blood to the head.
- I Chills in the room the whole day, especially after an evening nap.
- I Feeling of coldness all over, in the evening (without shuddering), without thirst (except at the commencement of the chilliness), and without being actually cold ; at the same time he feels out of humor, everything was disagreeable to him, even the meal ; two hours after he was covered with heat and sweat all over, without thirst.
- I Warmth all over, especially in the back.

Skin (Magnetis Polus Australis)

- I Corrosive itching in the evening, when in bed, on the back and other parts of the body.
- I Itching-stinging, tearing, here or there, in the evening, when in bed.

General Symptoms (Magnetis Polus Australis)

- I Bruised pain in all the limbs, so that he imagined he was lying on stones, on whatsoever side of the body he lay.
- I Cracking of the joints during motion.
- I Darting pains here and there, disappearing again immediately.

I Laziness and heaviness of the whole body, accompanied with a feeling of anxiety, as if he were threatened with paralysis, and as if he would fall, accompanied with a feeling of heat in the face and the whole body, mingled with shuddering.

I Lightness of the whole body.

I Stiffness of the joints.

o

MALANDRINUM (The Grease of Horses).

Mind (Malandrinum)

II Comprehension difficult.

II Memory weakened and impaired ; great difficulty in remembering what was read.

Stool (Malandrinum)

II Black, foul-smelling diarrhea, malaise and weariness.

II Dark brown, foul-smelling, almost involuntary diarrhea pains in abdomen.

II Dark brown, painless diarrhea.

II Diarrhea : acrid, yellow, offensive, followed by burning in anus and rectum.

II Diarrhea : yellow, bloody, slimy ; very changeable, worse in the morning ; acrid, excoriating ; child had a dried-up mummyfied appearance ; sleepless and has not nursed for 24 hours.

I Black, foul-smelling diarrhea ; weariness, nausea, dizziness.

I Dark, thin, cadaverous-smelling stool.

I Yellow, foul-smelling, almost involuntary diarrhea, and great weariness.

Back (Malandrinum)

II Intense pain across small of back.

II Pain along back as if beaten.

I Backache was intense in the sacral region ; in the dorsal region, under the shoulder blades, chiefly the left side ; it was almost unbearable. (Dr. B. from three doses of the 200th.)

Lower Limbs (Malandrinum)

II Deep rhagades, sore and bleeding, on soles of feet > in cold weather and after bathing.

II Soles of feet bathed in sweat, scald and burn when covered or warm.

I Feet "go to sleep" upon least provocation, a sensation never before observed.

I Large blisters on soles of both feet-no change of shoes skin exfoliated on both feet.

•

MEDORRHINUM.

Mind (Medorrhinum)

|| Dullness of memory and desire to procrastinate, because business seemed so lasting, or as if it never could be accomplished.

|| Fears he is going to die.

|| In conversation he would occasionally stop, and on resuming make remark that he could not think what word he wanted to use.

|| Is in a great hurry ; when doing anything is in such a hurry that she gets fatigued.

|| Time moves too slowly.

| Anticipates death.

| Cannot remember names ; has to ask name of her most intimate friend ; forgets her own.

| Cannot speak without crying.

| Cross through day, exhilarated at night, wants to play.

| Dazed feeling ; a far off sensation, as though things done today occurred a week ago.

| Dread of saying the wrong thing when she has headache.

| Entirely forgot what she had read, even previous line.

| Everything startles her, news coming to her seems to touch her heart before she hears it.

| Fear of the dark.

| Forgetfulness of names, later of words and initial letters.

| Great difficulty in stating her symptoms, loses herself and has to be asked over again.

| Great weakness of memory.

| Irritated at little things.

| Is always anticipating ; feels most matters sensitively before they occur and generally correctly.

| Loses constantly the thread of her talk.

| One night saw large people in room ; large rats running ; felt a delicate hand smoothing her head from front to back.

| Seems to herself to make wrong statements, because she does not know what to say next, begins all right but does not know how to finish ; weight on vertex, which seems to affect mind.

| Sensation as if all life was unreal, like a dream.

| Very impatient.

| Wild and desperate feeling, as of incipient insanity.

I Woke at an early hour with a frightened sensation, as if something dreadful had happened ; heavy weight and great heat in head ; could not rest in bed ; felt as if she must do something to rid her mind of this torture.

Head (Medorrhinum)

I Intense itching of scalp.

Eyes And Sight (Medorrhinum)

II Continuous watering of eyes, great heat and sensation of sand under lids.

I Aching in eyeballs, pressure and heat in vertex, a tendency to shut eyes.

I Decided tendency to irritation of edges of lids.

I Feeling as if she stared at everything, as if eyes protruded.

I Feeling of pain and irritation, and sensation of sticks in eyes, lids and especially inner canthi, redness and dryness of lids, congestion of sclerotic and sensation of a cool wind blowing in eyes, especially inner canthi.

I Neuralgic pain in eyeballs ; when pressing eyelids together ; < when rolling them.

I Ptosis of outer end of both upper lids, particularly left, requiring exertion to open them.

I Pulling pain in left lower lid from outer canthus ; could see lid twitch between these points.

I Swelling of upper lids ; soreness and smarting of edges.

Ears And Hearing (Medorrhinum)

II Child hard of hearing for six days.

I Partial or transient deafness ; pulsation in ears.

Nose And Smell (Medorrhinum)

II Epistaxis.

Face (Medorrhinum)

II Great pallor ; yellowness of face, particularly around eyes, as if occurring from a bruise (greenish yellow) ; yellow band across forehead close to hair.

II Neuralgia of r. upper and lower jaws, extending to temple.

II Rigidity of muscles of face, especially of lower lip, drawing it up tight to teeth ; jaws stiff, unable to open them ; deglutition nearly impossible ; throat filled with saliva.

II Tendency to stiffness in jaws and tongue.

Teeth And Gums (Medorrhinum)

II Hard swelling on r. upper jaw, as if in socket of a tooth gone since four years ; intense neuralgic pains extending to whole head, causing sleeplessness ; severe pains all over head, with external heat.

II Pale gums.

II Teeth have serrated edges, or are chalky and easily decay (Syphilinum ; teeth are cupped, decay at edges of gums and break off).

Mouth (Medorrhinum)

II Blisters on inner surface of lips and cheeks, skin peeling off in patches.

II Very sore mouth, ulcers on tongue and in buccal cavity, like blisters.

Appetite, Thirst, Desires, Aversions (Medorrhinum)

II Great craving for salt.

II Hiccough. Nausea ; with frontal headache ; after drinking water ; after dinner ; always after eating ; before eating.

II Violent retching and vomiting for forty-eight hours ; first glairy mucus, then frothy and watery, and lastly coffee grounds ; accompanied by intense headache, with great despondency and sensation of impending death ; during paroxysm was continually praying.

Scrobiculum And Stomach (Medorrhinum)

II Biliary colic with frequent vomiting and nausea ; diarrhetic stools, chilliness and perspiration on face and neck.

II Burning heat around the back, like a coal of fire.

II Congestion of liver.

II Cramps in stomach as from wind.

II Intense pain in stomach and upper abdomen, with a sensation of tightness.

II Sensation in pit of stomach as of a paper of pins that seemed to force themselves through flesh, causing her to rise and double up and scream ; pins seem to come from each side.

II Severe pain from abscess extending to r. shoulder and down elbow.

II Throbbing and thumping in region of suprarenal capsule, seeming to come from abscess or sore spot just below fifth rib, r. side ; creeping chills in region of r. kidney, throbbing, contracting, drawing and relaxing as if caused by icy cold insects with claws.

I Grasping pain in liver and spleen.

Abdomen (Medorrhinum)

II Ascites ; abdomen greatly distended ; palpitation showed water ; urine very scanty and high colored.

I Intense agonizing pain in solar plexus ; surface cold ; eructations tasting of sulphuretted hydrogen and, after eating, of ingesta ; applied r. hand to pit of stomach and l., to lumbar region.

Stool And Rectum (Medorrhinum)

II Baby, aet. 7 months, after summer complaint, great emaciation, diarrhea green, watery, slimy, yellow, curdled, like boiled potatoes chopped up with greens, thin, cream colored, watery, smelling like rotten eggs ; stools pass involuntarily ; apparently lifeless, except that it rotates head on pillow.

II Black stool.

II Can only pass stool by leaning very far back ; very painful, as if there was a lump on posterior surface of sphincter ; so painful as to cause tears.

II Child, aet. 15 months, brought on a pillow to clinic, apparently dead ; eyes glassy, set ; could not find pulse, but felt heart beat ; running from anus greenish yellow, thin, horribly offensive stool.

II Cholera infantum with opisthotonos, vomiting and watery diarrhea ; profuse discharge of blood and pus.

II Painful attacks of piles, not bleeding, hot swelling of left side of anus ; pin worms.

II White diarrhea.

I Constriction and inertia of bowels with ball-like stools.

Urinary Organs (Medorrhinum)

II Diabetic condition ; profuse and frequent urination.

II Nocturnal urination entirely ceased, the chamber empty in morning, a thing unknown for years.

II Pain in renal region, profuse urination relieves.

II Painful tenesmus of bladder and bowels when urinating.

II Passes an enormous quantity of high colored, strong smelling urine in bed every night ; thinks it is in afterpart of night, as he is always wet in the morning ; overwork and too much heat, or being in cold, aggravates this condition.

II Urine covered with a thick greasy pellicle.

II Very distinct bubbling sensation in r. kidney ; sensation of three bubbles in r. renal region, moving like bubbling in water, causing faintness ; deathly feeling in kidneys, with great depression of spirits, similar to effect of cold settling in renal region ; prostration after urination.

I Strong smelling urine.

I Urine high colored.

Male Sexual Organs (Medorrhinum)

II Burning in meatus during urination, and a feeling of soreness through whole urethra ; also after urinating, a feeling as if something more remained in urethra.

II Cannot retain urine more than an hour, after 5 or 6 P. M.

II Cannot retain urine through night.

II Gleet, gonorrhoeal discharge for twenty years ; urine stains clothes a dirty brown, discharges extremely slow, sometimes it takes half an hour to empty bladder, leaving him in a weak condition ; during urination, painful rectal tenesmus ; chilliness when bladder is too full, > by urination ; if he urinates after getting warm in bed, has to urinate every hour rest of night ; faint, indefinite sense of chilliness, followed by frequent calls to urinate, urine being hot, copious and followed by spinal chill and incontinence of urine on getting cold.

II Gonorrhoea for ten months ; during eight months mostly suppressed by drugs and injections ; for past two months flow persists, watery, transparent, but acrid and abundant, mixed with creamy liquid, stains linen yellow brown ; pain at end of penis during urination ; since third month of infection had heavy, drawing, wandering pains in r. arm, r. hip and left calf, < damp weather.

II Gonorrhoeal flow thin, transparent, mixed with opaque, whitish mucus, stains linen yellow.

II Impotence.

II Intense and frequent erections day and night.

II Pains along urethra while urinating, drawing burning.

II Profuse, yellow, purulent discharge from urethra, most copious in morning.

II Scanty, yellowish, gleet discharge, of many months' standing, showing most plainly in morning, gumming up orifice.

Female Sexual Organs (Medorrhinum)

II After very profuse menses, neuralgia in paroxysms in head, with twitching and drawing in limbs and cords of neck, which were like wires ; pain in lower abdomen, with profuse, yellowish leucorrhoea.

II Great sexual desire after menses, in a single woman.

II Menses very dark ; stains difficult to wash out.

II Ulceration of neck of uterus, which looked ragged and torn, inflamed and covered with stringy pus ; had gonorrhoea.

Voice and Larynx. Trachea and Bronchia (Medorrhinum)

II Bronchial catarrh spreading into larynx, swelling of tonsils and glands of throat extended also into ears, causing transient deafness.

II Choking caused by a weakness or spasm of epiglottis, could not tell which ; *larynx stopped so that no air could enter, only > by lying on face and protruding tongue.* (Acet. ac.) (Asthma : Dr. Miller, Glasgow.)

I Dryness of glottis, very annoying, with pain during deglutition ; great hoarseness.

Cough (Medorrhinum)

II Cough, < at night, causing retching.

II Cough, < on lying down, > lying on stomach.

Inner Chest And Lungs (Medorrhinum)

II Awful pains, in phthisis, in middle lobes.

II Constricted sensation at bottom of both lungs ; finally dull, heavy pain at top of left lung.

II Incipient consumption.

II Pain in r. shoulder as though it came from left, straight through.

Heart Pulse And Circulation (Medorrhinum)

II *Feeling of a cavity where heart ought to be.*

Neck And Back (Medorrhinum)

II Lumbago caused by straining in lifting.

II Pain in lumbar portion of spine ; myalgic ; induration of testes.

II Throbbing and thumping in region of r. suprarenal capsule seeming to come from abscess or sore spot just below fifth rib, r. side, under breast ; creeping chills in region of r. kidney, throbbing, contracting, drawing and relaxing ; as if caused by icy cold insects with claws.

I Heat in medulla and spine for a whole week.

I Pain straight through from left to r. shoulder.

Upper Limbs (Medorrhinum)

II Consumptive incurvation of nails.

II Pain commencing under left scapula, running down left arm to little finger, which pricked as if asleep.

II Rheumatic pain in r. shoulder and arm.

II Severe pain from abscess, extending to r. shoulder and down to elbow.

Lower Limbs (Medorrhinum)

II Cold and sweaty feet.

II Deformity of finger joints, large, puffy knuckles, swelling, stiffness and pain of both ankles ; great tenderness of heels and balls of feet ; the swellings of all affected joints were puffy like wind galls ; general condition < inland, > near shore.

II Old foot sweats, < during winter, for seven years.

II Pain like rheumatism along r. side, r. hip, left leg (upper, lower left) ; pains drawing, < in dampness ; left leg swollen near knee.

II Rheumatic pains in muscles of legs.

II Stiffness throughout body and joints.

II Sudden intense pain in left ankle, back of joint, on going to bed, could not move limb or body without screaming ; could find no position of comfort.

II Swelling and itching in soles, itching between toes, and pulling pains extending up to knees ; itching, painful, papulous eruption around waist, and hive like eruption wherever flesh is pressed on.

II Tenderness of soles so that he could not stand on them at all and had to walk on his knees.

I Almost entire loss of nervous force in legs and arms ; exhausted by slightest effort.

I Burning of feet, wants them uncovered and fanned.

I Kind of cramp in left calf at night, muscles knotted, > stretching ; not cramp, but knotting.

I Soreness in ball of foot under toes.

Nerves (Medorrhinum)

II Consumptive languor ; great general depression of vitality.

I Very tired.

Sleep (Medorrhinum)

II Becomes wide awake at 6 P. M., and continues so till 12, with entire passivity of brain and cessation of thought ; slight restlessness.

I Wakeful ; slept towards morning.

Fever (Medorrhinum)

II Chill came first at night, afterwards at various hours ; for instance, on two consecutive days at 2 P. M., then two days at 3, 4, 5 and 6 each, then at 7 P. where it remained for two weeks ; chill commenced in small of back, running up and down, lasted about an hour, and as it ceased, profuse and frequent urination appeared and continued during fever ; congestion of chest simulating pneumonia during fever, causing great alarm ; great renal distress during paroxysm ; thirst during fever for hot drinks ; fever continued for six or eight hours ; profuse sweat after fever ; great nervousness during paroxysm, was sure he would die ; intolerance of noise ; irritable.

II Chills for four months every day, commencing from 3 to 4 P. M. ; chill with headache, thirst, nausea, sometimes vomiting ; fever with headache, thirst, nausea, principally on head and neck ; pains from waist downwards ; frequent urination, dark color ; frequent, tasteless eructations ; constipation ; bad taste in mouth in morning.

II Profuse sweat about neck.

I Great burning heat all over body, with flashes of heat in face and neck.

Tissues (Medorrhinum)

I Obstinate rheumatism.

I Sequelae of acute articular rheumatism ; walks leaning on a cane, bent over ; muffled in wraps to ears, looking like a broken down man apparently soon to fall into his grave.

Skin (Medorrhinum)

II Copper colored spots (syphilitic) remaining after eruptions, turn yellow brown and detach in scales, leaving skin clear and free.

o

PSORINUM.

Mind (Psorinum)

II Anxious, full of fear, melancholic ; evil forebodings.

II Believe the stitches in heart will kill him if they do not cease.

II Depressed in spirits and hopeless.

II Driven to despair with excessive itching.

|| Feels the greatest anguish in head, with a whirling before eyes every day, from 5 A. M. until 5 P. M., since two years ; walks up and down his room wringing his hands and moaning continually, "Oh, such anguish ! Oh, such anguish !" only when he takes his meals he ceases moaning ; appetite is good.

|| Has been nervous about nine months ; was obliged to abandon all business ; has taken much quinine and other drugs ; a very disagreeable feeling about head ; mental depression ; thinks he will not recover ; has lost all hope ; cannot apply his mind to business ; confusion of senses, he cannot reckon ; attacks of numbness of legs and arms, left side< ; < on going to bed ; formication and crawling, with pricking and smarting on scalp, and some on extremities ; tongue white.

|| Intolerably self-willed, annoys those about him ; a boy, suffering from an eruption.

|| Melancholy after suppressed itch ; emaciated, pale, earthy complexion, weakness of limbs ; flushes of heat and palpitation prevent sleep ; sleep comes toward morning ; would like to stay in bed until midday ; aversion to work, indifference, weeping ; seeks solitude, despairs of recovery ; she is irritable and forgetful.

|| Much depression on account of an eruption on dorsum of hand which appears over night.

|| Very depressed, sad, suicidal thoughts.

I Anxiety, with oppression of chest.

I Despairing mood ; fears he will fail in business.

I Despairs of recovery ; thinks he will die ; hopeless ; especially after typhus, > from nosebleed.

I Discouragement.

I Great depression of mind during climaxis, with chronic abdominal disorders,

I Great fear of death ; anxiety about heart and dyspnea, with attacks of pain in chest.

I Greatest despondency, making his own life and that of those about him intolerable ; dry cough ; evening fever.

I His ideas are sad and joyless.

I Hypochondriasis, with hemorrhage from rectum.

I Irritable, peevish, passionate, noisy ; nervous, easily startled ; restless, hands tremble.

I Is so downhearted she could commit suicide, then is so full of phantasms.

I Melancholy, sorrowful, despairing.

I Religious melancholy. (Melilotus.)

I Restlessness : with eruption, in a child ; with oppression of chest.

I Sentimental : full of spleen, very low-spirited.

I She is very irritable, easily angered ; always thinks of dying.

I Vacillating, fearful ; mania.

I Very disagreeable mood ; impatient ; extremely ill-humored.

Sensorium (Psorinum)

I Vertigo ; mornings, objects seem to go around with him ; with headache ; eyes feel pressed outward ; with confusion and drawing in forehead, with roaring in ears.

Inner Head (Psorinum)

II Congestion of blood to head immediately after dinner.

II Congestion to head, cheeks and nose red and hot ; eruption on face reddens ; great anxiety every afternoon after dinner. Fifth month of pregnancy.

II Is always very hungry during headaches.

II Pain beginning over left eye and goes to r. ; < from hour to hour, then diarrhea and nausea, finally bloody vomiting ; dizziness, obliges her to lie down ; blur, and blue stars before eyes ; veins of temples much distended ; day before headache inordinate appetite ; also during first hours of pain ; < and brought on by change of weather, so that even in middle of night she is awakened by pain and always knows there has been a change ; soreness of stomach, sensitive to touch and pressure of clothes ; catamenia regular.

II Surging, drawing and digging in forehead with vertigo.

I Great congestion of brain, relieved nosebleed.

I Headache from repelled eruption.

I Headache preceded by : flickering before eyes ; dimness of sight or spots ; spectres ; objects dancing before eyes, black spots or rings.

Outer Head (Psorinum)

II A man, aet. 28, dark complexion, dark brown hair, had a spot on left frontal region, commencing at edge of hair and extending upward three-fourths of an inch ; the skin covering spot was many shades whiter than the surrounding skin, and the lock of hair growing on it had turned perfectly white ; after Psor. hair and spot became natural color.

II Moist, suppurating, fetid, also dry eruptions on scalp.

II Sensation as if head was separated from body.

II Tinea capitis et faciei.

I Crusta serpigiosa.

I Eruption on head, particularly on occiput, completely hiding scalp from view ; profuse exudation, soiling pillow, at night and causing excoriation of skin of nape of neck ; offensive smelling ; innumerable lice.

I Hair : dry, lustreless ; tangles easily ; glues together ; must comb it continually.

I Humid, scabby itching ; offensive smelling eruption on head, full of lice ; glandular swellings.

I Large humid blotches on head, with scabby eruptions on face.

I Profusely suppurating fetid eruption on head ; rawness and soreness behind ears.

I Pustules and boils on head, containing large quantities of pus ; severe itching, causing child to scratch so violently that blood flows ; formation of thick, dirty, yellow scabs, which when removed show a raw surface from which a yellow lymph exudes, which makes the linen stiff ; after removal new scabs form ; eruption

spreads on nape of neck, scalp and most of forehead ; the eruption is of very offensive odor ; such large quantity of lymph is exuded that head seems to stick to pillow ; child very restless, scratches head violently, and if prevented becomes irritable and screams ; large pustules on arms and body, which show no tendency to heal.

I Scurfy eruption of children ; large yellow vesicles around and between scabs.

Eyes And Sight (Psorinum)

II Great photophobia, walks with eyes bent upon ground ; scurfy eruption on face.

II Lids spasmodically closed ; intense photophobia and profuse flow of hot tears ; much pustular eruption on face ; large brown scab on r. eye, from beneath which pus pours forth abundantly when touched ; bowels costive ; appetite poor and only for dainties.

II Right eye red, internally and externally ; vesicles on cornea ; eruption on head.

I Acute ciliary blepharitis ; internal surface of lid chiefly affected ; photophobia ; strumous diathesis, with unhealthy, offensive discharges.

I Amaurosis, with scabby eruption on occiput and ears.

I Ciliary blepharitis r. to left, < morning and during day ; chronic cases ; subject to exacerbations.

I Confusedness before eyes after anxiety.

I Darkness before eyes and ringing in ears.

I Eyelids : swollen ; inflamed ; bloated ; child rubs eyes ; puffy ; greatly swollen, closely pressed together ; thickened ; tendency to styes ; itching, especially in canthi ; herpetic eruption ; scrofulous inflammation, covered with thick crusts, whole body covered with branlike tetter.

I Eyes become gummy.

I Fiery sparks before eyes.

I Heat, redness and pressure in eyes ; tendency to catarrhal inflammation ; lids slightly agglutinated during morning.

I Inflammation of eyes with burning.

I Inflammation of lids, internal surface much congested ; great photophobia, cannot open eyes, lies on face.

I Lachrymation.

I Objects seem to tremble for a few moments and get dark.

I Ophthalmia, with pressing pains, as if sand were in eyes.

I Photophobia : when walking in open air ; with inflammation of lids.

I Recurrent pustular inflammation of cornea and conjunctiva ; chronic form ; scrofulous basis.

I Rheumatic, chronic and blepharophthalmia.

I Scrofulous inflammation of eyes ; ulceration of cornea.

I Serous choroiditis ; some ciliary congestion and great haziness of vitreous, so that optic nerve was only discerned with great difficulty and then was found decidedly hyperemic, as was the whole fundus ; headache, especially in morning ; constant profuse sweating of palms of hands.

I Vision blurred ; 'black spots before eyes ; flickering ; dancing about of objects.

Hearing And Ears (Psorinum)

II Otorrhea : with headache ; thin, ichorous and horribly offensive, like rotten meat ; very offensive, purulent (watery, stinking diarrhea) ; brown, offensive, from left ear, for almost four years ; chronic cases following scarlet fever.

II Right ear a mass of crusts and pus, the crusts extended behind auricle to occiput upward upon parietal bone nearly to vertex, forward to r. ear and over cheek ; upon edge of region involved small vesicles filled with clear fluid, which became yellow, then crusted, and pus flowed from beneath crusts.

II Scurfs on ears, and humid scurfs behind ear.

I Discharge of reddish earwax or fetid pus.

I External ears raw, red, oozing, scabby ; sore pain behind ears.

I Herpes from temples over ears to cheeks ; at times throws off innumerable scales ; at others shows painful rhagades with yellow discharge, forming scurfs ; fetid humor ; itching intolerable.

I Itching in ears ; child can hardly be kept from picking or boring in meatus.

I Meatus externus scabby.

I Pustules : on and behind concha ; behind left ear.

I Singing, cracking, humming, buzzing and ringing in ears with hardness of hearing.

I Ulceration of membrana tympani.

Nose And Smell (Psorinum)

II Bloody, purulent discharge from nose.

II Chronic catarrh ; dropping from posterior nares, so as to awaken him at night ; hawking quantities of lumpy mucus gave temporary relief from feeling of fullness ; mucus in nose would dry like white of egg, needed to be forcibly removed.

II Dry coryza with stoppage of nose.

II Pain in liver, < from sneezing.

II Septum narium inflamed, with white, suppurating pustules.

I Acne rosacea.

Face (Psorinum)

II An offensive-smelling, crusty eruption extending over whole face for three months, had completely closed eyes.

II Eruption on face of a child ; whole face covered with a crust, lips and eyelids swollen, aversion to light, large moistening spots on head and behind ears.

II Moist scab behind ears with dry tetter on back of head, on both cheeks extending upward to eyes and downward to corners of mouth, reddish, very closely packed, millet-seed like, itching, dry pimples, with frequent loose stools ; a child one and a half years old.

II Pain as if lame in condyle of jaw.

II Painful tension and pressure in r. zygoma, towards ear.

II Sweat of face with general heat.

I Closely packed, itching pimples on both cheeks from eyes to corners of mouth.

I Coppery eruption on face.

I Corners of mouth sore, often ulcerated ; sycotic condylomata.

I Crusta lactea on face and scalp, especially over either ear and cheek, exfoliating numerous scabs, or it cracks and discharges a yellow, fetid humor.

I Humid eruptions on face ; whole face covered with humid scurfs or crusts, with swelling of lips and eyelids and humid soreness behind ears.

I Lips : painful ; swollen, particularly upper ; dry ; burning ; brown and black, dry ; ulcerated ; swollen and covered with scurfs.

I Much roughness of skin of face ; eruption on forehead between eyes ; stools very offensive.

I Scabby face ; especially cheeks from ears ; lips and eyelids swollen, sore about eyes.

I Tinea faciei.

I Ulcers in face.

Teeth And Gums (Psorinum)

II Looseness of teeth ; they feel so loose, fears they may fall out, < from touch, especially from teeth.

I Gums : ulcerated, bleeding.

I Toothache : < at night and from cold ; > from warmth.

Taste And Tongue (Psorinum)

I Loss of taste with coryza.

I Taste : bitter, goes off when eating or drinking ; foul, much mucus in mouth ; > in fresh air ; bitter with yellow-coated tongue ; flat, sticky, dinner tastes oily.

I Tongue : dry, tip feels burnt as far as middle, he has hardly any taste ; tip very dry, as if burnt, painful ; white, yellow ; thickly coated with whitish-yellow slime ; ulcerated.

Mouth (Psorinum)

I Dryness of mouth ; burning.

I Ulcers in mouth.

Throat (Psorinum)

- || Cutting tearing pain in throat on swallowing.
- || Sensation of a plug or lump in throat impeding hawking.
- || Steam arising from fat causes immediate constriction of throat and chest.
- || Tension and swollen feeling in throat.
- || Throat burns, feels scalded.
- | Accumulation of mucus in throat and mouth.
- | Difficult swallowing, throat feels swollen.
- | Dryness ; scraping sensation in throat.
- | Dryness in throat with thirstlessness.

Appetite, Thirst. Desires, aversions (Psorinum)

- || Thirst : during dinner ; with dryness of throat ; especially for beer, mouth feels so dry.
- | Diminished appetite : after typhus ; but great thirst ; during convalescence.

Eating and Drinking (Psorinum)

- || Immediately after dinner, congestion of blood to head.
- || Pain in chest extending to shoulder, < after cold drinks.
- || Waterbrash when lying down, > on getting up.
- | Drinking causes cough.

I Eructations : sour, rancid ; tasting and smelling like rotten eggs ; room is filled with an offensive odor. Arn., Graph., Ant. t. Arn. : especially in A. M. ; Ant. t. : at night ; Graph. in A. M. only, after rising, disappearing on rinsing the mouth.

I Nausea : with poor appetite ; in morning ; with backache, after suppressed itch ; morning with pain in small of back ; all day, with vomiting.

Scrobiculum and Stomach (Psorinum)

- || Frequent oppression of stomach, especially after eating.
- || Gastric bilious affections.
- || Weakness of stomach.
- | Dyspepsia ; eructations, flatus and stools like spoiled eggs.

Hypochondria (Psorinum)

|| Deep-seated stitching, pressing pain in region of liver, < from external pressure and lying on r. side ; pain hinders sneezing, laughing, yawning, coughing, deep inspiration and walking.

I Chronic hepatitis.

Abdomen (Psorinum)

II Colic : removed by eating ; > passing fetid flatus.

II Flatulency with disorders of liver.

I Abdominal affections during climaxis, with a high degree of ill-humor.

I Bloating abdomen.

I Chronic abdominal affections.

I Constant feeling of emptiness and looseness in abdomen ; sensation as if intestines were hanging down.

I Inguinal hernia ; hernial sac infiltrated.

I Pains in abdomen after eating ; flatulency and tendency to diarrhea ; > when flatus passes.

Stool and Rectum (Psorinum)

II Boy, aet. four months, whitish bad-smelling diarrhea ; constantly crying, with drawing up of knees as if in pain ; rattling cough, cries all night ; acts as if he had earache in left ear, from which there was a slight discharge ; seldom urinates ; after second dose, sixteen hours after first, broke out all over his head and face, with a small pimply eruption, a vesicle at the apex of each pimple, which exuded lymph which dried in a thick brown scab, which gradually fell off in a day or two, and in two weeks left the skin perfectly clean.

II Hemorrhage from rectum ; in old women large quantities of blood discharged at once, with constipation and hypochondriasis.

II *Horribly offensive*, nearly painless, almost involuntary, dark and watery stool ; only at night and most towards morning.

II *Involuntary stool* during sleep.

II Semi liquid, brownish, indelible, insufferably nasty ; passed during sleep, at 1 and 4 A. M., with undigested food. Infantile diarrhea.

II Sensitive hemorrhoidal pain in rectum.

I Before stool : griping pains about navel.

I Cases which do not respond promptly to the indicated remedy, the children having dirty, yellow, greasy skin, with a partially developed eruption on forehead and chest, with constant fretting and worrying.

I Cholera infantum ; stools dark brown, watery, of an intolerably offensive odor ; liquid, mucous or bloody and excessively fetid.

I Cholera.

I Chronic diarrhea ; offensive stools.

I Diarrhea : after severe acute disease ; at night ; early in morning ; when rising in morning ; in childbed ; when weather changes (general condition).

I Diarrhea, preceded by colic.

I Frequent thin stool, with eruption on head.

I Lienteria.

I Obstinate constipation : with coryza and obstruction of nose ; pain in small of back ; blood from rectum ; due to torpor of rectum ; lasting three or four days, due to inactivity of rectum ; stool on third or fourth day was accompanied by severe pains which induced patient to withhold effort as much as possible.

I Soft stool : passed with difficulty ; from weakness.

I Stool : dark brown, very fluid and foul-smelling ; having the smell of rotten eggs ; mostly in children in their first or second summer ; green mucus, or bloody mucus ; smells like carrion, < at night ; frequent, liquid ; involuntary ; nearly painless.

I Unpleasant burning in rectum.

Urinary Organs (Psorinum)

II Involuntary urine, cannot hold it ; vesical paresis. Typhus.

II Scanty urination nearly every half hour, with burning in urethra and in condylomata.

I Enuresis : wets bed at night ; again during full moon ; obstinate cases.

I Urine : dark brown, with reddish sediment ; loaded with pus ; frequent, scanty ; burning and cutting in urethra ; thick whitish ; turbid ; red deposit ; cuticle forms on surface ; profuse.

Male Sexual Organs (Psorinum)

II After suppressed gonorrhoea ; rheumatism, lameness ; conjunctivitis, with granulations ; intense photophobia ; pain darting around, through head : other eye sensitive to light.

II Boy, aet. 7, suffering since birth ; r.-sided inguinal hernia, about three inches of intestine descending to testicle through widely opened inguinal canal ; upon applying a bandage severe inflammation of tunica vaginalis occurred, which yielded to Pulsat., but returned every time bandage was applied ; child gradually grew miserable and thin and lost all appetite ; fever set in, and a large amount of water collected in tunica vaginalis ; as the case improved a painful, burning, itching excoriation with acrid discharge appeared upon inner surface of prepuce and upon corona glandis.

II Gleet of twelve years' duration.

II Hydrocele, caused by repeated inflammation, in consequence of pressure from a truss.

II Seven large, moist, itching, occasionally burning condylomata on prepuce ; every night nocturnal enuresis ; during day must urinate nearly every half hour ; urination scanty and accompanied by burning in urethra and condylomata ; lips ulcerated, particularly at corners of mouth ; in several localities, but particularly in popliteal spaces, dry, herpetic eruption, not itching.

II Sycotic excrescence on edges of prepuce, with itching and burning.

I Absence of erections ; parts flabby, torpid.

I Aversion to coition ; impotence ; want of emission during coitus.

I Chronic painless discharge from urethra, leaving yellow stain upon linen.

I Hydrocele.

Female Sexual Organs (Psorinum)

II Dysmenorrhoea near climaxis.

II Knotty lump above r. groin ; even a bandage hurts.

II Left ovary indurated after a violent knock ; followed by itching eruption on body and face.

II Menstrual disorders during climaxis.

II Metrorrhagia.

II Leucorrhoea, large lumps, unbearable in odor ; violent pains in sacrum and r. loin ; great debility.

I Amenorrhoea : in psoric subjects when tetter is covered by thick scurfs ; with phthisis.

I Ulcers of the labia.

Parturition. Pregnancy. Lactation (Psorinum)

II Mammae swollen, painful ; redness of nipples, burning around them.

II Pimples itching violently, about nipples ; oozing a fluid. Second month of pregnancy.

I During pregnancy : fetus moves too violently ; abdomen tympanitic ; nausea ; vomiting ; obstinate cases.

Voice and Larynx. Trachea and Bronchia (Psorinum)

II For eleven years hay fever, coming on about 20th of August.

II Voice weak, trembling.

I Hoarseness ; when talking, phlegm sticks in larynx.

I Suffocative and crawling sensation in larynx, producing a paroxysmal, dry, hacking cough.

I Talking is very fatiguing.

Respiration (Psorinum)

II Convalescents go out for a walk, instead of being invigorated return home in order to get breath or to lie down so they can breathe more easily, feel < instead of > from being in open air.

I Anxious dyspnea, with palpitation and pain in cardiac region.

I Asthma, as if he would die ; precursor of hydrothorax.

I Chest expands with great difficulty ; cannot get breath.

I *Dyspnea* : < when sitting zip to write, > when lying down, congestion to head after dinner, great despondency ; < the nearer arms are brought to body.

I Short breath or want of breath.

Cough (Psorinum)

II Cough aggravates pain in liver and pain in chest extending to shoulder.

II Cough with expectoration ; asthma, thinks he will die.

II Cough, causing tearing from centre of chest to throat, all on r. side ; cough < at night ; urine escapes when coughing.

II Dry cough, pain in the chest for last three months, a constricting pressure at fourth and fifth ribs near sternum, excessive irritability and ill-humor.

II Severe, dry cough with oppression of chest and pain as if everything in chest were raw and scratched ; fever in evening ; great depression of spirits, making life burdensome to him.

I Chronic blennorrhoea of lungs, threatening phthisis.

I Cough : from tickling in larynx ; dry, hard, caused by tickling in trachea ; in evening with pains in chest and throat, passing off when she is quiet ; produced by talking ; with sensation of weakness in chest ; dry, with sensation of heaviness in chest ; dry, with soreness under sternum, with stitches in chest ; dry, with constricting pain in chest ; < mornings when awaking and evenings on lying down ; coughs a long time before expectorating ; periodic attacks ; chronic, spasmodic ; of twenty-five years' duration.

I Cough with expectoration of green mucus, nearly like matter ; especially in morning when waking and in evening when lying down, with nausea ; it sticks firmly and he can only expectorate with difficulty.

I Drinking causes cough.

I Expectoration : of blood with hot sensation in chest ; yellowish-green.

Inner Chest and Lungs (Psorinum)

II Chest pains from coughing.

II Constriction of chest when inhaling steam from fat.

II Dull pressure in r. side of chest, extending thence over whole chest, < bending forward in writing, not by motion or deep inspiration ; dry cough with expectoration of small lumps of mucus ; speaking affects him very much ; great prostration after preaching, so that he must rest a long time to recuperate ; voice is not husky, but it requires all his strength to get through with his work ; chest narrow, shoulders projecting. Phthisis.

II Fixed pain in r. side of chest.

II Hot sensation in chest.

II Pain in chest comes by fits ; great anxiety ; a feeling of ulceration under sternum ; chest inflates only with much exertion ; coughs a long time before beginning to expectorate. Phthisis pulmonalis.

II Pain in chest, as if raw, as from subcutaneous ulceration.

II Pains in chest grow more severe two or three times a day, begin with chilliness and trembling, followed by heat one hour in duration ; great anxiety of heart and mind with, fear of death, dyspnea and restlessness ; attacks pass off with sour, clammy sweat and chilliness ; sweat however occurs every night independent of attack.

II Stitches : in sternum, with backache ; from behind forward in chest and back when breathing ; in r. side of chest when breathing ; in chest (left side).

II Tedious recovery in pneumonia.

I Chronic blennorrhoea of lungs.

I Hydrothorax.

I Suppuration of lungs. Phthisis pulmonalis.

Heart, Pulse and Circulation (Psorinum)

II Gurgling (gluckern) in region of heart, particularly noticeable when lying.

II Rheumatic pericarditis ; pulse 144 ; skin dry ; pain in head and limbs, but more particularly in shoulder ; dyspnea, with pain in region of heart ; effusion, indistinct heart sounds ; bellows murmur with first sound ; inability to lie down.

II Stenosis of left ostium venosum ; purring in region of apex ; cyanotic lips ; dyspnea and shortness of breath when walking in open air ; > lying down.

II Stitches in cardiac region, low gurgling extending towards heart, for a moment breathing is impossible.

I Dyspnea : with palpitation ; with pain in cardiac region.

I Pain in heart > when lying down, thinks the stitches will kill him if they continue.

I Palpitation : with anxiety ; mental disquietude, dislike for work ; from coughing ; in those suffering from hepatic disorders.

I Pericarditis of psoric origin ; > lying quietly.

Neck and Back (Psorinum)

II Backache when walking, with stitches in sternum.

II Backache with constipation.

II Constant pressing pain in small of back, < from motion.

II Herpetic eruption on side of neck extending from cheek.

II Nape of neck excoriated by discharge from eczema capitis.

I Backache after suppressed eruption.

I Excessive backache.

I Severe backache, as if bruised, cannot straighten out.

I Spina bifida.

Upper Limbs (Psorinum)

II Copper-colored eruption or red blisters on backs of hands.

II Dry tetter on wrists with rheumatism in limbs.

II Herpes in palms of hands ; itching tetter.

II Itch like eruption on wrists with tearing in limbs.

II Nails brittle.

|| Swelling and tension of backs of hands and of fingers.

|| Tetter on arm with small millet-like eruption exuding a yellow fluid ; itches intensely in heat.

| Eruption in bends of elbows and around wrists.

| Itching between fingers ; vesicles.

| Malignant boil ; on hand a cone-shaped scab the size of a quarter of a dollar on a base as large again, bluish-red and strongly demarcated, where scab extends over ring there is another moist, white ring which forms a new scab ; much tension and burning.

| Pustules on hands, near finger-ends suppurating.

| Sweaty palms, especially at night.

| Tearing in arms.

| Trembling of hands.

Lower Limbs (Psorinum)

|| Carries ; rachitis.

|| Chronic rheumatism in limbs, with a dry eruption on wrists.

|| Corns between second and third toes of left foot.

|| Dry herpes, especially in bend of knees.

|| Eruption about joints makes walking difficult, as if encased in armor.

|| For four or five weeks, feeling when walking as if left foot were pulled around inward ; < for last two weeks, so that he sometimes looked to see if it were really so. Locomotor ataxia.

|| For many weeks gouty pains, etc. ; dry cough ; constrictive pressure and cutting, tearing pain at sternum near fourth and fifth ribs ; greatest despondency and ill-humor.

|| Gout in lower extremities.

|| Hands and feet feel as if broken early in morning and after a little work.

|| Hands moist, with cold, clammy sweat, the very touch of which was unpleasant ; profuse sweating of feet ; feet very painful, causing shuffling gait.

|| Ischias : sciatica.

|| Knees give way under him.

|| Large swelling about ankle.

|| Old itch eruption on inner side of thigh and in popliteal space.

|| Purpura on inner side of thigh.

|| Vesicles becoming ulcers, on feet.

I Arthritis : rheumatism, especially in chronic forms.

I Chronic gonitis.

I Heat in hands and feet.

I Herpetic and itching eruption especially in bends of joints, in bends of elbows and in popliteal spaces.

I Oozing blisters on legs, from small pustules, increasing in size, with tearing pains.,

I Pain in knee caused by a fall a year ago.

I Paralysis of legs from suppression of eruption on arms.

I Tearing in limbs ; in left knee and left axilla.

I Ulcers : on legs usually about tibiae and ankles or other joints ; ulcers are indolent, slow to heal ; on lower legs with intolerable itching over whole body ; on feet.

Nerves (Psorinum)

II A man, aet. 21, was obliged to run until nearly exhausted ; although strong and well before, he now became weak, perspiring easily and had severe pain in r. side, < by coughing, laughing, motion.

II Debility : independent of any organic disease ; loss of appetite ; tendency to perspiration on exertion and at night ; after acute diseases ; after protracted diseases or loss of fluids ; after typhus, with despair of recovery ; thinks he is very ill when he is not ; appetite will not return ; in the evil consequence of suppressed itch, especially after large doses of sulphur.

II Subsultus tendinum.

II Trembling and chilliness, with attacks of pain in chest.

II Very weak and miserable after suppressed itch.

I Constantly increasing debility, with abdominal affections.

I Constantly tired and sleepy ; very little labor exhausts him ; exhausted after riding in a wagon.

I Loss of strength, with cough, with oppression of chest.

I Malaise : feels tired out.

I Nervous, restless, easily startled.

Sleep (Psorinum)

II Child apparently well, but at night would twist and turn and fret from bedtime till morning and next day be as lively as ever.

II In morning lies in same position as when he fell asleep.

I Sick babies will not sleep at night, but worry and fret and cry. (See Jalapa.)

I Sleepless after 12 P. M., from congestion to head.

I Sleepy by day ; sleepless at night, from intolerable itching ; dyspnea ; congestion to head.

Fever (Psorinum)

II After ague color of face worse.

II Profuse sweating relieves all the complaints ; chronic diseases. (Calad.)

II When taking a walk profuse sweat with consequent debility, taking cold easily.

I Chilliness in evening on upper arms and thighs, with thirst ; drinking causes cough, then heat and cough, with oppression of chest ; and trembling, with attack of pain in chest.

I Heat : at night and dryness in mouth ; in afternoon or evening, feels as if he would lose his senses, with thirst ; in evening, with delirium, great thirst, followed by profuse sweat ; when riding in a carriage ; sudden over whole body, with trickling perspiration all over face.

I Internal shivering, creeping chills and icy-cold feet.

I Profuse night sweats of phthisis.

I Sweats easily, weak.

I Typhus : picks bedclothes, reaches for objects in air ; profound debility.

I Want of perspiration ; dry skin.

Skin (Psorinum)

II A girl, aet. 18, sallow complexion but cleanly appearance ; constant irritation of the different parts of body, day and night, compelling her to scratch ; numerous pediculi corporis cling to neck, back and shoulders ; menstruation had never occurred ; anxious, depressed and tearful ; inner angles of eyes filled with gummy mucus, heat and itching in them, in evening ; corners of mouth sore.

II Burning itching pustule after vaccination.

II Eczema behind ears, on scalp and in bends of elbows and armpits, accompanied by abscesses affecting bones ; nothing relieved, but the eruption disappeared, to reappear again, years after, on wrists ; there was then a patch on each wrist as large as a half dollar, with intense itching, preventing sleep, with constant desire to scratch.

II Itch : dry on arms and chest, most severe on finger joints ; followed by boils ; inveterate cases with symptoms of tuberculosis ; in recent cases, with eruptions in bends of elbows and around wrists ; repeated outbreak of single pustules after main eruption seems all gone.

II Large suppurating pustules on hands, particularly near ends of fingers ; had eight or ten within a few weeks, itching of body particularly while in bed.

II Moist, itching condylomata.

II Psoriasis ; psoriasis syphilitica.

II Pustules or boils on head, particularly on scalp ; scalp had a dirty look and emitted an offensive odor ; fine, red eruption, forming small white scales ; pustules on hands.

II Retrocession of eruption. Measles.

II Scaly condition of skin of whole body ; skin has a dirty, tawny color, although carefully kept ; much itching causing desire to scratch, which gave but temporary relief ; some months back instep showed signs of

eruption, which soon became a thick, dirty-looking mass of scales and pus, painful and violently itching ; at times pain kept her awake at night ; disease of at least a dozen years' standing.

II Severe pains in limbs and dark, burning spots, so that skin of whole body except face resembled that of a mulatto ; itch had been suppressed five times.

II Skin has a dirty, dingy look, as if patient never washed ; in some places looks coarse as if bathed in oil ; sebaceous glands secrete in excess ; body always smells dirty.

II Suppressed eruptions.

I Abnormal tendency to receive skin diseases.

I After suppressed itch : urticaria in attacks, after every exertion ; tuberculosis ; single pustules often appear.

I Consequences of itch suppressed by sulphur ointment.

I Crusta serpigiosa.

I Crusty eruptions all over body.

I Eruptions bleed easily and constantly tend to suppurate.

I Herpes : with itching and burning ; with biting-itching, with meal-dust, humid.

I Herpetic eruptions.

I Itching : when body becomes warm ; and stinging in many parts at same time ; intolerable, < in bed and from warmth ; scratches until it bleeds ; over whole body, when rubbed, small papules and vesicles arise : between fingers ; in knee joints ; in bends of knees ; terrible, of whole body at night, preventing sleep.

I Moist herpes after suppressed scabies ; intolerable itching when getting warm ; < before midnight and in open air.

I Pemphigus.

I Skin dirty, greasy looking, with yellow blotches here and there.

I Skin inactive ; want of perspiration.

I Ulcers : deep, penetrating, ichorous ; on face and legs ; old, with fetid pus ; violently itching ; scrofulous, with swelling of bones.

o

SECALE CORNUTUM.

Mind (Secale cornutum)

II Laughs, claps her hands over her head, seems beside herself.

II Sensation of intoxication while undressing, to retire (third day and for several successive evenings).

Sensorium (Secale Cornutum)

II Sensation of intoxication while undressing.

Outer Head (Secale Cornutum)

II Scalp sore.

Eyes and Sight (Secale Cornutum)

II Exophthalmos with struma.

II Pustulous conjunctivitis and blepharitis.

II Retinitis diabetica.

II Suppuration of cornea ; < from warm applications.

I Cataracta senilis.

Nose and Smell (Secale Cornutum)

II Nose stopped up on left side as with a solid plug.

II Nosebleed : blood dark, runs continuously, with great prostration, small, threadlike pulse ; in old people or drunkards ; of young women ; from debility.

Teeth and Gums (Secale Cornutum)

II Difficult dentition ; great weakness ; vomiting of everything taken ; great thirst ; pale face ; eyes dim, sunken ; dry heat, with rapid pulse ; restlessness and sleeplessness.

Throat (Secale Cornutum)

II Severely paralyzed both in swallowing and in speaking ; could scarcely take food without great danger of choking ; speech reduced to a whisper ; could not bear heat or covering and would throw all covering off. Post-diphtheritic paralysis.

II Throat sore on left side running up into ear.

I Diphtheria : loss of strength ; rapid loss of sensibility ; numbness of extremities ; painful tingling and crawling on tongue ; dry gangrene ; apathy ; dilated pupils ; burning pains of affected parts ; stammering speech ; absence of all reaction.

I Follicular pharyngitis ; hawking up of little follicular exudates.

Scrobiculum and Stomach (Secale Cornutum)

II Attacks of severe pressure and constriction in region of stomach extending through to spine, extremely painful and followed in half an hour by vomiting of tasteless fluid or of contents of stomach, thereupon an intermission of several hours occurred ; during attack region of stomach felt as if contracted, and on percussion gave a tympanitic note ; has three to four attacks daily.

II Bilious vomiting, with cramping pains in stomach ; burning in stomach extending up esophagus ; head sunk upon breast, face pale, yellowish, voice weak, pulse small. Cardialgia.

II Hematemesis ; attacks preceded by pains in epigastrium and nausea, pain going to left side when pressure is made in epigastrium ; marked protrusion in left hypochondrium, with pain ; blood red, never containing particles of food, and when collected in basin appears more like bloody serum than pure blood and is of offensive odor, quantity vomited not very large ; frequent chilliness at night, followed by profuse sweat ; strength not much impaired ; appetite and sleep good.

I Hemorrhage from stomach ; lies still with great weakness but no pain ; face, lips, tongue and hands deadly pale ; skin covered with cold sweat, pulse frequent, threadlike ; oppression ; abdomen soft, without pain. Hematemesis.

I Severe anxiety and burning at pit of stomach.

Abdomen (Secale Cornutum)

II Aneurism of mesenteric artery, in women.

I Pain in abdomen with burning in stomach.

I Pains in hypogastric region.

Stool and Rectum (Secale Cornutum)

II Aversion to heat or being covered, with icy coldness of extremities.

II Colliquative diarrhea.

II Diarrhea : very exhausting ; pernicious ; very offensive ; involuntary, profuse, watery, putrid, brown ; discharged with great force ; very exhausting ; urine suppressed ; painful with great prostration ; painless with tingling and numbness in limbs ; putrid, fetid, colliquative, patient does not want to be covered or to be near the heat, but prefers to be in the air or wishes to be fanned ; sudden attacks ; of children, discharges whitish, watery ; chronic in overfed children, great prostration ; during August ; great stools undigested, or watery, at times yellowish, also greenish, with forcible expulsion, accompanied by discharge of flatus ; paralytic weakness of sphincter ani with involuntary discharges.

II Five to ten minutes after taking least quantity of food, severe colic which made her bend double and cry out ; pain begins between region of stomach and navel, extends thence to sides and rest of abdomen and down to sacral region, accompanied by severe urging and tenesmus, followed by a thin, slimy, yellowish stool with some relief of pain ; four to five such attacks follow each other ; then relief until she eats again ; four to five evacuations during night ; she compares pains to labor pains ; great thirst ; thick mucous coating on tongue ; sleep disturbed ; prostration. Diarrhea.

II Patient cold, almost pulseless, with spasmodic twitching of muscles in various parts of body ; spreads fingers asunder ; eyes sunken, features pinched ; much spasmodic retching although not much vomiting ; skin harsh, shrivelled, dry, as if there were no moisture left in system ; urine suppressed ; tingling or formication all over body ; stools profuse, watery, ejected with great violence ; is cold but cannot bear to be covered. Cholera.

II Profuse prostrating evacuations, severe painful cramps in feet, toes, hands and fingers which are spread apart or extended toward back of hands ; cramping pressure in stomach ; dry, wrinkled, cold skin ; cyanotic color. Cholera.

II Stools yellowish-white, slimy, undigested, escaping involuntarily, < at break of day.

II Uncomfortable fullness of abdomen, with transient pinching pains in upper abdomen as from flatus ; at night severe cutting pains throughout whole abdomen ; restless anxious tossing about, with short and unrefreshing naps ; during night anus firmly closed, "as if locked up ;" in morning frequent short watery evacuations, in gushes, preceded by cutting pains in abdomen.

II Vertigo, cramps or drawing in calves of legs, rumbling in abdomen, nausea, stools in rapid succession, brownish or colorless, rapid prostration, coldness of limbs, tongue but slightly coated.

I Cholera Asiatica, with collapse, sunken, distorted face, particularly mouth ; crawling sensation as from ants.

I Cholera infantum ; cholera morbus ; cholera Asiatica.

I Cholera infantum ; profuse undigested stools, watery and very offensive, discharged by fits and starts and followed by intense prostration ; pale face, sunken eyes, dry heat, quick pulse, restlessness and sleeplessness ; great aversion to heat and to being covered.

I Cholera with more retching than vomiting.

I Constipation.

I Diarrhea after cholera.

I Hemorrhage from bowels.

I Interminable diarrhea in summer, which resists everything, especially in scrofulous children ; putrid, fetid and colliquative ; choleraic symptoms, with cold, clammy perspiration ; sinking spells at 3 A. M. (not the restless anguish of Arsenicum).

Urinary Organs (Secale Cornutum)

II Hematuria in a boy suffering from suppuration of the glands of neck after scarlet fever ; urine also very albuminous ; anasarca ; great thirst.

I Bloody, albuminous urine.

I Enuresis : old people ; pale, watery or bloody urine.

I Paralysis of bladder.

I Passive hemorrhage ; blood thin ; blood corpuscles wanting in consequence or dissolution ; or painless discharge of thick black blood in consequence of kidney disease ; coldness of body ; cold perspiration on forehead ; great weakness. Hematuria.

Male Sexual Organs (Secale Cornutum)

I Weak memory after exhausting coition ; impotence.

Female Sexual Organs (Secale cornutum)

II A woman, aet. 45, passed through a normal confinement seven years ago ; miscarried about two years ago, hemorrhage kept up for five months under allopathic treatment, with exacerbation during time of menstruation ; after ceasing for seven months, hemorrhage set in again with slight intermission of one or two weeks ; excessive anemia ; sunken features ; skin cold and dry ; pulse small and quick ; heavily coated tongue ; loss of appetite ; headache ; since five days, daily ten or twelve painless stools, of mucous, watery, sometimes foul smelling masses ; thin, black, foul-smelling bloody discharges.

II An excessive menstrual flow every two weeks, lasting seven to nine days ; for last four weeks flow is continuous ; very weak and thin ; has severe pains in loins and uterine region ; bearing down pains as if in labor.

II Feeble and extremely emaciated, skin flaccid, face very pale and sunken, with an expression of suffering, mucous membranes pale and cool, hands and feet deficient of natural warmth, action of heart quickened, breathing short and oppressed, pulse very small, 120 ; abdomen distended ; os uteri very open, with indented and puffy edges, flaccid and soft, vagina tender and cool ; manual examination caused much uneasiness and flooding ; violent headache limited to one spot, throbbing in temples, roaring in ears, giddiness on slightest movement ; enfeebled nervous system showed extraordinary excitability ; many times in day, and especially at night, cramps in calves and spasmodic twitching of limbs, causing exhaustion, remains several hours in bed as if paralyzed ; digestion and sleep disturbed to some extent ; hemorrhage still continued, even in horizontal position, and elevation of pelvis caused no diminution in large quantity of blackened coagula which were constantly passing, while least movement increased discharge in a very great degree. Chronic passive hemorrhage.

- II Gangrene of whole vaginal mucous membrane ; on holding apart the labia this membrane was found of a dark slate .color, emitting the characteristic odor.
- II Menses irregular ; every four weeks for three to four days, copious dark-red fluid discharge of blood, with pressing, labor like pains in abdomen ; constipation ; pressure in occiput ; afterward continuous discharge of watery blood, until next period.
- II Menstrual colic ; pains so severe as to frequently cause spasms ; uterine region very sensitive to touch ; high fever ; pains > when flow appears.
- II Menstrual colic ; pale face ; coldness of limbs ; cold sweat ; small, suppressed pulse ; tearing, cutting pains in abdomen.
- II Metritis ; tendency to putrescence ; inflammation caused by suppression of lochia or menses ; discharge of thin black blood, a kind of sanies, with tingling in legs and great debility. Metritis.
- II Metrorrhagia ; relaxed condition of body ; depressed, anxious state of mind ; unusual drowsiness by day ; gush of thin black blood on least movement of body ; general feeling of prostration ; diminished temperature of body ; wooden, numb. feeling in lower extremities.
- II Pain in ovaries and uterus.
- II Putrescence of uterus ; abdomen distended, not very painful ; discharge from vagina, brownish, offensive ; ulcers on external genitals discolored and rapidly spreading ; burning hot fever, interrupted by shaking chills ; small, sometimes intermittent pulse ; great anguish, pain in pit of stomach, vomiting decomposed matter ; offensive diarrhea ; suppressed secretion of urine ; skin covered with petechial and miliary eruptions or shows discolored, inflamed places, with a tendency to mortification ; the patient lies either in quiet delirium or grows wild with great anxiety and a constant desire to get out of bed. Metritis.
- II Since last delivery menses too seldom and very irregular, last time rather copious in consequence of unusual exertion ; at night dreamed she was ascending the stairs with a heavy load, and suddenly a clot of blood came away and the blood seemed to gush forth ; the alarm awakened her and she found that she was bleeding fast ; on following morning strength greatly reduced ; lips and whole body, even tongue, deathly pale ; pulse could not be distinguished ; frequent fainting fits ; periodic pains with expulsion of clots of blood and between these attacks constant oozing of thin bright blood.
- II Uterus about an inch from labia inferiora, membrane around it felt hard, while rest of mucous membrane of vagina was very much relaxed and gathered into a fold at lower part.
- I Cancer and gangrene of uterus.
- I Discharge from vagina almost black, fluid and very fetid.
- I Dreadful bearing down, dragging out feeling in lower abdomen, so that her life is almost unbearable ; every four or five days profuse, thick, yellow discharge from vagina ; hesitation in urinating ; rheumatism. Hysteralgia.
- I Incessant metrorrhagia.
- I Leucorrhoea : in thin, scrawny women, with prolapsus uteri ; green, brown, offensive ; like cream, from weakness and venous congestion.
- I Menses : too profuse and lasting too long ; with tearing :and cutting colic, cold extremities, cold sweat, great weakness and small pulse ; or with violent spasms.
- I Painless flooding in feeble, cachectic, dyscratic women, or such as have long resided in tropical climates.
- I Partial prolapsus of uterus for eight months after a forceps delivery ; dysuria ; sense of weight over pubes as if contents of abdomen would fall forward.

I Prolapsus of three months' standing ; frequent severe cutting pains in abdomen ; occasional nausea.

I Uterine hemorrhage : when uterus is engorged ; with pains in sacrum, extending down thighs and pressing into lower abdomen of a pregnant woman ; profuse protracted flow ; tearing, cutting colic, cold extremities and cold sweat, weak, hemorrhage < from slightest motion, blood thin and black, black, lumpy or brown fluid, of disgusting smell ; black liquid blood.

I Uterine hemorrhage, flow passive, dark and may be offensive ; tingling or formication all over body, 'holds her fingers spread asunder, asks to have her limbs rubbed ; finally lies. unconscious and cold.

I Uterine ulcers feel as if burnt, discharge putrid, bloody fluid.

Pregnancy. Parturition. Lactation (Secale cornutum)

II After labor, pale, weak ; uterus distended, burning pains therein, hard, painful to least touch ; discharge of black, coagulated or brown, watery offensive-smelling blood ; throbbing, tearing pains in thighs extending down to toes ; pain < from motion ; strong pulsations in umbilical region, which could be felt by the hand ; pulse at wrist weak and rapid ; frequent yawning.

II After lifting a heavy weight during sixth month of pregnancy severe pains in stomach, abdomen and small of back and a pushing-down sensation ; violent movements of fetus ; cold feet ; numbness and tingling in feet ; small, weak pulse.

II After-pains : too long and too painful.

II Cessation of lochia, with fever ; inflammation of uterus, subsequently an abscess opened through vagina.

II During eighth month of pregnancy violent convulsions with frothing at mouth, etc., followed by variable spasms ; insensibility and clonic spasms, < at every pain ; on return of consciousness complained of dull frontal and occipital headache and incessant uterine pains. Premature labor.

II During labor : prolonged bearing-down and forcing pain in uterus ; pains irregular ; pains too weak ; pains feeble, or ceasing ; everything seems loose and open, no action ; fainting fits.

II During pregnancy : frequent and prolonged forcing pains, particularly in thin, ill-conditioned women ; cramps in calves.

II Extremely violent pains, almost without intermission, she seemed to be in the last stage of labor, but on examination os was found about the size of a half dollar, thick and somewhat rigid.

II Extremely violent pressing labor pains, os, however, being only about as large as a ten cent piece ; hysterical convulsions.

II Lochia : dark, very offensive ; scanty or profuse ; painless or accompanied by prolonged bearing-down pain ; suppressed, followed by metritis ; suddenly changes character and becomes of a dirty brown or chocolate color, with fetid odor, grows sad and melancholy and fears death ; of too long duration.

II Post-partum hemorrhage, with relaxation of uterus, only temporarily relieved by compression : after-pains excessive, < when child nursed.

II Prone to abortion in third month ; had passed through five, although she kept her bed as soon as pregnant ; some labor pains with bloody discharge ; was able to attend to her household duties and went to full term.

II Violent after-pains with hemorrhage arising from irregular contractions ; the longitudinal fibres alone contracting in such a manner as to leave a sulcus in middle, making it appear as if uterus were split open from top to bottom.

II When advanced about seven and a half months in pregnancy was taken with labor pains, wriggling and not distinctly intermitting ; os tincae open, and about size of a shilling ; dullness and slight aching of head ; despondent.

II While the head was passing in lower strait, she was suddenly seized with violent convulsions lasting about three or four minutes, followed by a stupid state with stertorous breathing and uneasy moaning as if from pain.

I A sensation of constant tonic pressure in uterine region ; causes great distress ; desires fresh air ; does not like to be covered.

I After abortus : difficult contraction of uterus ; thin, black, foul-smelling discharge.

I Fever with frequent watery stools. Puerperal fever.

I Hour-glass contraction.

I In women who are much exhausted from venous hemorrhage ; thin, scrawny women ; the breasts do not properly fill with milk.

I Labor ceases, and twitchings or convulsions begin.

I Lack of milk with much stinging in mammae.

I Puerperal convulsions with opisthotonos.

I Retained placenta, after miscarriage, especially when occurring during early months of pregnancy ; offensive discharges ; patient cold and often almost pulseless from loss of blood ; uterine contractions very imperfect, or else prolonged tonic contraction.

I Retained placenta, with constant, strong bearing-down in abdomen, or with relaxed feeling of parts.

I Strength of uterus weakened by too early or perverted efforts.

I Strong tendency to putrescence ; discharge of sanious blood, with tingling in legs and great prostration ; urine suppressed ; offensive diarrhea ; voice hollow with difficult breathing, feeble and inaudible ; burning fever interrupted by shaking chills, does not care to be covered ; cold limbs ; cold sweat over whole body ; gangrene.

I Suppression of milk ; the milk will not flow from the breast.

I Thin, scrawny women, skin shrivelled, dry and harsh, sallow face, weak in labor ; pains seem to be entirely wanting ; uterus flabby ; bearing down in sacral region, a sort of prolonged urging feeling in abdomen.

I Threatened abortion : more especially at third month : with copious flow of black, liquid blood ; false labor pains, with bloody discharge ; in feeble, cachectic women, having a wan, anxious countenance, pulse almost extinct, fear of death ; convulsive movements.

I Uterine pains prolonged but ineffectual.

Cough (Secale Cornutum)

II Concussive cough ; profuse perspiration ; sleepless nights ; inclination to colic ; diarrhea ; bloatedness of abdomen ; emphysema. Bronchitis.

I Spitting of blood, with or without cough.

Inner Chest and Lungs (Secale cornutum)

II Expectoration of dark, frothy, rather viscid blood, brought up by a slight cough and amounting to a teacupful in four hours ; a spot as large as a crown piece on r. side of chest to r. of nipple, dull on percussion, with bronchial respiration and mucous râle over that part.

Heart, Pulse and Circulation (Secale Cornutum)

II Palpitation ; hot forehead ; inclination to sleep ; spasmodic shocks from r. half of chest into r. arm and leg ; in paroxysms every two or three hours ; oftener in night, after each meal ; less in open air ; coldness and numbness of r. hand, with tingling in fourth and fifth fingers ; loss of muscular power and feeling in hand ; after sexual excess.

I Palpitation of heart : with contracted and frequently intermitting pulse.

I Pulse : often unchanged even during violent attacks ; generally slow and contracted, at times intermittent or suppressed ; somewhat accelerated during heat ; small ; empty, weak ; threadlike, in hemorrhages.

Neck and back.(secale cornutum)

II Hard, hoarse cough, with but little expectoration ; pains nearly all over front part of chest, < from coughing and motion ; for several years tenderness of lower cervical and upper dorsal spinous processes, with stiffness of neck ; < from every exertion or strain upon spine ; pressure upon diseased portion of spine produces pain there, as well as all through chest, with irritation to cough.

II Stitches in upper dorsal vertebrae (between shoulders), constant when sitting, intermittent when standing, at times extending into hands, < by pressure upon vertebrae ; frequent formication through all limbs ; at times rigidity and spasmodic stretching of fingers so that for several minutes he cannot sew ; frequent pressure and swelling beneath epigastrium ; pain in back < when sewing.

I Myelitis diffusa.

Upper Limbs (Secale Cornutum)

II Left thumb spasmodically drawn toward dorsum of hand, followed in a few minutes by cramping and flexure of rest of fingers toward palmar surface ; hands feel numb like velvet ; next day both hands became affected and after several days felt a tingling and stitches in legs, followed by heaviness of same, < after walking, generally appearing while sitting, while cramps in hands always appear after using them.

I Contraction of fingers.

I Gangrene of fingers ; senile gangrene.

I Numbness and insensibility of hands and arms.

Lower Limbs (Secale Cornutum)

II Beginning senile gangrene ; swelling and livid coloring of r. foot extending to malleoli ; foot cold ; severe indescribable pains.

II Cold gangrene of limbs.

II Cramp in calves.

II Dry gangrene of foot with constant severe, burning tearing pains.

II Gangrene of toes.

II Profuse, stinking and corrosive perspiration of feet, softening and bleaching soles and destroying quickly stockings and shoes ; existing two months in a girl eighteen years old.

|| Senile gangrene commenced in great toe of r. foot and slowly extended ; foot was livid and swollen ; all the symptoms pointed to its complete loss.

I Burning in feet.

I Cramps in calves of legs and soles of feet, disturbing sleep at night and hindering walking in pregnant women.

I Cramps in hand and toes.

I Fuzzy feeling in limbs.

I Paraplegia.

I The feet seem asleep and stiff.

I Toes of r. foot spasmodically drawn upwards, continuously during day and occasionally at night, causing a peculiar limping gait ; this cramp was accompanied by no pain, but by a very tiresome sensation rendering walking, particularly going up and down stairs, very difficult ; tendons running along dorsum of foot to toes were tense as wires and the corresponding muscles of leg larger and harder than normal ; now and then slight sensation of coldness in back and also a peculiar buzzing (formication) in spine.

I True anthrax, rapidly changing into gangrene.

Nerves (*Secale cornutum*)

|| After abortus, spasms with full consciousness, afterward great exhaustion ; heaviness in head and tingling in legs.

|| Burning : in all parts of the body as if sparks of fire were falling on them.

|| Collapse from choleroïd diseases, etc., with cold skin, yet unable to bear warmth.

|| Complains of great weakness, constipation, heaviness in epigastrium, formication in legs and cramps ; at night while asleep gets epileptiform attacks of which she knows nothing next morning except that she feels greatly prostrated and has a constant heaviness in head.

|| Convulsions first occurring after a fright when a little girl, returned after each confinement.

|| Convulsions.

|| Expression of countenance varied every moment from a constant play of the muscles ; eyes rolled about, pupils dilated, tongue jerked out, head moved about from side to side ; arms in constant action with most diverse movements ; snatched objects rather than took hold of them, could hold nothing securely ; trunk also in constant motion ; urine could not be retained ; pulse small, weak, quick ; heart beat tumultuously ; appetite poor ; bowels torpid ; aching in occiput ; sensation of formication in extremities ; memory impaired ; speech difficult, hurried ; no relief at night ; staggered about the house almost all night .

|| Numbness of extremities ; paralysis of some parts ; painful tingling (like crawling of ants) on tongue.

|| Paralysis of lower extremities, in a woman past climacteric ; a hard, sensitive tumor in one of her breasts had been developing for several years, but for the last year or two had rapidly increased in size and become very painful ; by continued application for several months of a yellowish salve the tumor was enucleated ; in about a month after ceasing to use the salve, a peculiar torpor or deadness was felt in great toes, which extended to whole foot and ankle joint ; feet seemed large and heavy and could be moved only by moving whole limb ; gait shuffling as if feet were dragged along by lifting legs ; slight numbness in hands.

II Restlessness ; extreme debility and prostration.

II Sensation of soreness in abdomen ; formation of large lumps and swellings in abdomen ; r. hand very weak, particularly fingers, so that she could hold nothing, nor sew with that hand ; when placing open hand to side and taking it away a spasm of hand occurs, the fingers are spread apart and she cannot close the hand ; rapid alternations of heat and cold in hands and feet ; cramps in legs ; icy coldness of knees ; trembling of r. arm and hand while eating, must use the left ; sensation of coldness in stomach, > for a short time by warm drinks ; habitual constipation.

II Sinking spells from diarrhea, at 3 A. M.

II Suddenly fell to ground, but without loss of consciousness ; on every attempt to walk the knees would sink from under him ; especially r. ; while sitting cannot bend forward or backward without losing his equilibrium ; arms slightly weak, sensation of touch not affected ; general dullness ; difficult speech ; inclination to weep ; complains of headache and pain in lumbar region ; sleep poor ; urine and feces escape involuntarily ; no stool for several days ; at times oppression of breathing.

II The muscular twitchings usually commence in face and spread thence all over body, sometimes increase to dancing and jumping.

II Twitching of single muscles ; twisting of head to and fro ; contortions of hands and feet ; labored and anxious respiration.

I Chorea associated with menstrual irregularities.

I Hyperesthesia of cutaneous nerves, especially of spine.

I Neuralgia, caused by pressure of distended veins upon a nerve trunk.

I Paralysis after spasm, and apoplexy, with rapid emaciation of affected parts and involuntary discharges from bowels and bladder.

I Paralysis with rapid emaciation, with relaxation of sphincters.

Fever (Secale Cornutum)

II Aversion to heat or to being covered ; may feel cold but does not wish to be covered.

II Exhausting perspiration, accompanied by evening fever and alarming cough.

II Heat with thirst and hot skin.

II Intense, icy coldness of skin, with shivering ; pale, sunken face.

II Violent shaking followed by violent heat, with anxiety, delirium and almost unquenchable thirst.

Kept limbs drawn up close to body, are too cold to lie straightened out ; they ache but are too cold to straighten out.

I Burning heat, interrupted by shaking chills, then internal burning heat, with great thirst.

I Chill with thirst.

I Cold limbs, cold sweat, great weakness.

I Cold surface ; sunken pale face and blue lips ; will not be covered ; tingling in limbs ; holds hands with fingers widely spread apart ; cold, clammy sweat ; speech feeble, stuttering. Ague.

I Great tendency to typhoid. Intermittent.

I Severe and long-lasting dry heat, with great restlessness and violent thirst.

I Sweat : all over body, except face ; profuse cold, cold limbs ; from head to pit of stomach ; especially on upper body ; cold, clammy over whole body ; colliquative.

I Violent chill of but short duration ; followed soon after by internal burning heat, with great thirst.

Tissues (Secale Cornutum)

II Collapse from choleroïd diseases.

II Gangrene : from anemia ; external injuries, application of leeches or mustard ; > from cold, < from heat ; dry, of old people.

It destroys the activity of the cord ; convulsive twitchings and shocks, painful contractions, tetanic manifestations ; perfect paralysis, with increased reflex activity ; most excruciating spinal pains, especially in sacral region ; paralysis of bladder and rectum ; tendency to gangrene ; rapid emaciation.

I Dry gangrene of extremities, parts are dry, cold, hard and insensible, of a uniform black color and free from fetor ; large ecchymoses, blood blisters on extremities, becoming gangrenous, black suppurating blisters ; limbs become pale, cold, shrivelled or lead-colored, losing all sensibility.

I Lymphatic tumors.

I Neuralgia caused by pressure on nerves by a distended vein.

I Rapid emaciation of paralyzed parts.

I Ulcers : bleeding ; becoming black ; feeling as if burnt ; painless ; pricking, producing a prurient sensation ; pus putrid ; < at night, touch, from external warmth ; > from cold.

Skin (Secale Cornutum)

II Boils, small, painful, with green contents, mature very slowly and heal in same manner ; very debilitating.

II Formication : with a sense as if mice were creeping under skin ; on face, gums and other parts of body ; in extremities with tearing, stinging pains ; in tips of fingers, lasting several weeks, with a partial loss of sensibility ; over whole body.

II Purpura haemorrhagica.

I Bloody blisters on extremities, becoming gangrenous.

I Cachectic females, with rough skin ; pustules showing tendency to gangrene.

I Carbuncles ; extensive ecchymoses.

I Crawling : all over body ; between skin and flesh ; and jerking under skin.

I Indolent ulcer, ichorous, offensive pus, > from cold. General desquamation in scarlatina.

I Petechia and miliary eruptions.

I Ulcers turn black, copious vomiting of a mixture of a thick, black, pitchy, bilious or shiny matter.

I Varicose ulcers and enlarged veins in old people.

I Variola pustules of abnormal appearance, either fill with a bloody serum or dry up too soon.

o

SYPHILINUM.

Mind (Syphilinum)

II Cross, irritable, peevish.

II Terrible dread of night, not on account of cough so much as on account of mental and physical exhaustion when she awakes ; it is intolerable, death is preferable ; she fears to prepare for night and is positively in abject fear of suffering, in form of exhaustion on awaking ; it is < by cough, but it is quite independent of cough as she wakes in this awful state ; always < as night approaches ; leaves her about daylight, which she prays for. Spring cough. Had a dose of Syphilinum cm., twenty-four hours before ; on third night her anxiety and cough returned. though very much less. Another dose of Syphilinum cm., next morning, removed all symptoms, including aphonia, and exhaustion requiring brandy.

II Very despondent, does not think he will ever get better.

Head (Syphilinum)

II Coronal headache.

II Dirty eruption on scalp.

II Lancinating pain in occiput, < invariably at night, and causing sleeplessness, but always ceasing with the coming light of morning.

II Nervous chills preceded by aching pains in head, especially in occipital and integuments thereof, head feeling heavy, sore, congested ; also frontal headache about one-half or two-thirds inches wide across forehead under eyebrows ; aching pains below waist, in pelvis and extremities, especially in tibia, which is sensitive to touch ; pains commence about 4 P. M., culminate about midnight in delirium, and cease entirely at daylight.

II Syphilitic cephalalgia in occiput, intolerable, extending to nervous ganglia of neck, causing hardening of cords ; attacks at irregular intervals, especially after excitement.

II Syphilitic headache for many months, piercing, pressing excruciating over r. eye ; extending deep into brain ; losing continuity of thought and memory ; makes repeated mistakes in figures.

I Headache : linear, from or near one eye backward ; lateral ; frontal ; from temple to temple ; deep into brain from vertex ; as from pressure on vertex ; in either temple, extending into or from eye > by warmth ; in bones of head ; < by heat of sun ; after sunstroke.

I Headache through temples, thence vertically like an inverted letter T.

I Headaches accompanied by great restlessness, sleeplessness and general nervous erethism.

I Neuralgic headache causing sleeplessness or delirium at night, always commencing about 4 P. M. ; < at from 10 to 11 P. M., and ceasing at daylight.

Eyes (Syphilinum)

II Acute left conjunctivitis, with considerable pain in eyeball, photophobia and lachrymation, followed by iritis ; nocturnal aching in eyeball, pain extremely violent from 2 to 5 A. M. ; sight impaired.

|| At 1 P. M. scalding lachrymation of r. eye with shooting therein, followed by shooting from around eye into eye ; eye red and closed ; this lasted about an hour, then decreased, ceasing about 3 P. m. ; recurred for two successive days and again four days afterwards in a slighter degree, but at same hour.

|| Both eyes glued in morning ; conjunctiva injected ; photophobia, constantly wears a shade.

|| Eyes dull.

|| Eyes swollen and closed with syphilitic ophthalmia, pus running out of them.

|| Eyes very red and inflamed.

|| L. eyeball covered with fungus-like growth, pain intense < at night.

|| Left eye closed, upper lid swollen as large as half an English walnut ; deep red, not much pain, with oozing of purulent matter from between lids.

|| Neuralgia every night, beginning about 8 or 9 P. M., gradually increasing in severity until it reached its height about 3 or 4 A. .d., and after continuing thus for two or three hours, gradually decreased and finally ceased about 10 A. M. ; attacks gradually get more severe and last longer ; first feels cold all over, almost a shiver ; then soreness as if beaten in r. half of head, extending a little beyond middle line on vertex ; in about thirty minutes scalding lachrymation from r. eye, with shooting backward therein ; eye is very red and closes, with photophobia ; gnawing pains extend down r. side of face and whole of nose ; head is worst when eye is bad ; during paroxysm r. eye feels as if lids were open wide, and cold air blowing on exposed eye ; she perceives a horizontal band across pupil of r. eye hindering sight ; this came on soon after paroxysms commenced ; eye > by placing handkerchief on head and letting it hang over eyes, also by gentle pressure, though she cannot bear much pressure ; it is more painful when lying on r. (affected) side when also r. side of head feels sore ; r. eye red, and red vessels run all over it, converging towards iris ; r. pupil horizontally oval ; r. iris looks dull and there is a slight brown hue around pupil ; left eye normal ; attacks seem to have originated from sitting at a window in a cold draft, r. eye being next window.

|| On turning eye to left feels momentary coldness in inner half of r. eye.

|| On waking, gum in r. canthus.

|| On walking across room, r. eye sensitive to air, aches on using it.

|| Ophthalmic pains, < at night, > by cold water.

|| Pain in r. inner canthus as if blood went there and could go no farther, also in r. temple.

|| Pain in r. inner canthus as if blood went there and could go no further, also in r. temple.

|| R. eye alone affected, congestion of conjunctiva and sclerotica, with some chemosis ; lids inflamed, esp. at outer canthus ; sensation of sand in eyes ; lids agglutinated in morning ; great photophobia (hereditary syphilis).

|| Red papulous eruption round left inner canthus, with isolated pimples on side of nose, cheek and eyebrow ; these pimples were red, with depressed centre, circumscribed areola, became confluent where they were most dense ; pimples bleed when scabs come off ; agglutination of lids.

|| Redness and swelling of outer half of both lower tarsal edges.

|| Syphilitic iritis, intense pain steadily increasing night after night ; < between 2 and 5 A. M., coming almost at the minute and ceasing same way.

| Acute ophthalmia neonatorum.

I Acute ophthalmia neonatorum.

I Chronic recurrent phlyctenular inflammation of cornea ; successive crops of phlyctenules and abrasion of epithelial layer of cornea ; intense photophobia ; profuse lachrymation ; redness and pain well marked ; delicate, scrofulous children, especially if any trace of hereditary syphilis remains.

I Chronic recurrent phlyctenular inflammation of cornea ; successive crops of phlyctenules and abrasion of epithelial layer of cornea ; intense photophobia ; profuse lachrymation ; redness and pain well marked ; delicate scrofulous children, especially if any trace of hereditary syphilis remains.

I During sleep lids adhere ; in infantile syphilis.

I Paralysis of superior oblique.

I Upper lids swollen.

Ears (Syphilinum)

II Calcareous deposit on tympanum.

II Catarrhal or nerve deafness with marked cachexia.

II Complete deafness ; nothing abnormal to be seen.

II Deafness gradually increasing until she could scarcely hear at all.

II Gathering in left ear which discharges a great quantity of pus (hereditary syphilis in a child).

II Intense earache in r. ear, incisive pains thrusting into ear ; purulent watery discharge from ear with pain.

II Small, acrid, watery discharge occasionally from ears, no deafness (ozaena).

Nose (Syphilinum)

II L. side of nose inside and out very sore, likewise lips and chin ; sores itching and scabbing over ; after Syphilinum lm. much better in twelve hours, many drying up and scabs falling off, leaving skin beneath of a dull-reddish copper color.

II Offensive, thick yellow-green nasal discharge ; during sleep dry scabs form in both nostrils ; following an application of salve for sore eyes ; left submaxillary gland, which had been swollen and indurated, softens, discharges, and, after forty-five days, begins to heal slowly.

II Ozaena syphilitica. (Syph. brought out an eruption of sores with a fiery-red base on nose over frontal sinuses.)

I Attacks of fluent coryza.

I Itching in nostrils.

Face (Syphilinum)

II A boy, aged 20 months, fretful, peevish, cross and crying, tossing in his sleep, grinding his teeth, face dotted with papules filled with a watery yellowish matter, most on edges of lids ; teeth irregular, arms and legs emaciated, very tottery on his feet, very nervous.

II Face pale.

II Facial paralysis r. side., thick speech, hemicrania, jactitation of r. eye and lid.

II Lips and teeth covered with bloody mucus.

II Nose and cheeks covered with eruptions and scabs in layers rising to a point.

II Spasmodic twitching of many muscles, esp. in faces (**paralysis agitans**), with great melancholy and depression of spirits.

I Dark purple lines between ala : nasi and cheeks.

I Itching, scabby, eczematous eruptions singly or in clusters, looking like herpes.

Teeth (**Syphilinum**)

II Felt like a worm in tooth, could not tell which tooth.

Mouth (**Syphilinum**)

II Aphasia, difficulty of finding words ; debility.

II Fetid breath.

II Putrid taste in mouth before epileptic fit.

II Syphilitic destruction of hard and soft palates.

II Tongue very red and thick ; covered with herpetic eruption, two deep cracks running lengthwise on each side of median line, making it difficult to swallow.

II Twenty ulcers in mouth, every part involved, top and under side of tongue, lips, buccal cavity, fauces and nose.

I Herpetic eruption in mouth, tonsils, hard palate and fauces, completely covering inside of mouth and throat, making it difficult to swallow even liquids.

I Tongue coated white, edges indented by teeth.

Throat (**Syphilinum**)

II Chancrous ulcer extending across velum palati to left pillar of pharynx, which was congested and thickened, interfering very much with his speech ; voice husky.

II Chronic hypertrophy of tonsils.

I Acute pharyngitis.

Appetite (**Syphilinum**)

II Appetite indifferent and capricious.

II Total loss of appetite for months, little or nothing satisfies him ; formerly was generally ravenous.

I Aversion to meat.

I Dyspepsia ; flatulence, belching of wind ; nervous dyspepsia.

I Loss of appetite.

I Tendency to heavy drinking ; alcoholism.

Stomach (Syphilinum)

II Vomiting for weeks or months due to erosion from superficial ulceration of lining of viscus, herpetic, of syphilitic origin.

I Heartburn with pain and rawness from stomach to throat pit, often with cough.

Abdomen (Syphilinum)

II Large painless bubo in r. groin opened and discharged freely.

I Inguinal bubo.

Stool and Anus (Syphilinum)

II Bilious diarrhea at seashore, painless, driving her out of bed about 5 A. M. ; stools during day ; later causing excoriation ; face red, suffers from heat ; occasional painless, whitish diarrhea when at home, > always by going to mountains.

II Bowels torpid for five weeks.

II Fissures in anus and rectum.

II Lower portion of rectum hanging out like a ruffle, looking like a full-blown rose fully three inches in diameter, and sensitive ; constant weak dragging sensation in rectum, extending as far as sacrum.

II Obstinate constipation for many years ; rectum seemed tied up with strictures, when injections were given agony of passage was like child-bearing.

II Stools too light-colored.

II Stools very dark and offensive.

II Two indurated ulcers at mouth of anus somewhat sore ; slight itching of anus.

I Chronic constipation, with fetid breath, earthy complexion, gaunt appearance.

I Obstinate cases of cholera infantum.

Urinary Organs (Syphilinum)

II Frequent urging to urinate all night, at least from 7 P. M. until 5 A. M., or sunset to sunrise.

II Profuse urination after chill ; passed during night nearly a chamberful.

II Rich lemon-yellow scanty urine.

II Urination difficult and very slow ; no pain, but a want of power, so that he has to strain.

I Urine infrequent, not oftener than once in twenty-four hours, scanty, of a golden-yellow color ; after Syphilinum lm., woke next morning with great distention of abdomen and pain in region of kidneys ; rising she passed a large quantity of normal colored urine, after which the distention and pain were relieved ; next day regular urination, watery.

Male Sexual Organs (Syphilinum)

II After suppressed chancre, disease attacked testes and scrotum, which became painful and swollen ; this was supposed to be cured, but ever since, every few weeks, if exposed to damp weather would be seized with pain as if in kidneys, seemingly traversing ureters, but instead of passing into bladder followed spermatic cord, down groins and into testes ; pain agonizing, chiefly in cord, in present attack in r. ; pricking in chancre, as though punctured by pins.

II Chancroid, phagedenic, spreading rapidly ; buboes commencing in each groin.

II Constant pain in anterior part of r. thigh, < while standing, painful all night, preventing sleep ; bubo in left inguinal region size of a pigeon's egg, purple, fluctuating ; night sweats.

I Buboes.

I Chancre on penis, third in two years, all on same spot.

I Chancre on prepuce.

I Inflammation and induration of spermatic cord.

Female Sexual Organs (Syphilinum)

II Acrid discharge causing violent itching and inflammation of external organs < at night from warmth of bed, parts very tender ; itching and inflammation > during menses.

II Nocturnal < of r. ovarian pain, preventing sleep.

II Painful menstruation.

II Sore on r. labium majus, extending to left

II Soreness of genitals, and muco-purulent discharge, in a child.

II Uterine and all surrounding parts loose, soft and flabby ; profuse, thick, yellow leucorrhoea ; constant pain across small of back.

I Ovaries congested and inflamed ; tendency to ovarian tumors.

I Profuse yellow leucorrhoea < at night ; in sickly, nervous children.

I Uterine and ovarian diseases with pronounced nervous disorders, esp. in married women.

I Yellow offensive leucorrhoea, watery or not, so profuse it daily soaks through napkins and runs to heels of stockings if much on her feet.

Respiratory Organs (Syphilinum)

II Attacks of spasmodic bronchial asthma for twenty-five years ; they come on only at night after lying down or during a thunder-storm, producing most intense nervous insomnia, entirely preventing sleep for days and nights.

I Diseased cartilages of larynx.

I Violent attacks of dyspnea, wheezing and rattling of mucus, from 1 to 4 A. M.

Cough (Syphilinum)

II Cannot lie on r. side, as it causes a dry cough.

II Dry, racking cough, with thick, purulent expectoration, caused by a sensation of rasping or scraping in throat, always < at night.

II Dry, sharp, hacking cough without expectoration, but with rawness, scraping and burning from fauces to stomach pit ; with a whoop in inspiration and a choking sensation from fauces to bifurcation of bronchia, great mental distress.

II Expectoration without cough, quite clear, white, feels like a round ball and rushes into mouth.

II Muco-purulent expectoration, greyish, greenish, greenish yellow, tasteless.

II Whooping-cough with terrible vomiting.

Chest (Syphilinum)

II Chronic asthma ; in Summer, especially when weather was warm and damp ; most frequently in evening, passing off at daybreak ; soreness of chest, with great anguish and inability to retain a recumbent position ; in Winter severe bronchial cough succeeded by asthmatic attacks ; a regular type of chills and fever developed ; suffered from this many years before. Within 24 hours after taking the 1m. the character of the asthma changed, during the night it disappeared, and the hour which had been that of relief, now became that of aggravation. The attacks only lasted a few minutes, and gradually becoming less severe, did not return for several months.

II Rattling in chest and throat.

I Angina ; ptosis left eye ; facial paralysis left side, slight aphasia ; impotence (relieved).

I Eczematous herpetic eruptions on chest.

I Pain and pressure behind sternum.

Neck And Back (Syphilinum)

II Caries of cervical spine with great curvature in same region, directly forward, occiput sinking down to a level with it and resting on protuberance of curvature ; often nearly a teaspoonful of calcareous matter would be discharged at a time and on evaporating it a quantity of dry powder, looking like phosphate of lime, would be left ; pain in curvature always < at night (no proof of syphilis).

II Caries of dorsal vertebrae with acute curvature, numerous cloacae communicating with diseased bone, one much larger than the rest, exuding a sanious, offensive pus, and surrounded with proud flesh ; great thickening and induration of surrounding parts from effusion of lymph ; percussion or pressure not endurable ; two abscesses in groins, left having been opened a year before r. about a month ago ; least motion gave him great pain by day ; and terrific pain by night ; for five months, every night most intense neuralgic pains, commencing generally from 5 to 7 P. M. and never terminating till about daylight or about 5 A. M. ; pains in muscles of loins, generally in left, sharp, cutting spasms, terrible to bear, preventing sleep and forcing him to cry out ; < at least motion, and slightly > by warm poultices.

II Enlargement of cervical glands and a number of pedunculated pin-head warts on neck ; cured by *syco-syphilinum*.

II Enlargement of glands in different parts of the body, particularly abundant about the neck ; indurated and slightly painful causing a sensation of uncomfortable fullness and suffusion in face, throat and head.

II Enormous swelling of glands in head and neck ; no relief from any remedy. Cured with *Buboin syphilitica* cm.

II Great pain in back in region of kidneys, < after urinating.

II Nocturnal aggravation of pains in back, hips and thighs.

II Psoas abscess first left then r., latter discharged more than a quart of offensive greenish pus when opened ; severe nocturnal pains, affecting upper sacral, lower dorsal and left cervico facial regions, steadily increasing ; they occurred twice, each time twenty-one days after either psoas abscess had been opened.

I Heavy aching and stiffness from base of neck up through muscles and cords into brain.

Upper Limbs (Syphilinum)

II Fingers and thumbs have runarounds (infantile syphilis).

II Hands badly ulcerated on backs.

II R. second finger is swollen and stiffened.

I Rheumatism of shoulder-joint or at insertion of deltoid, < from raising arm laterally.

Lower Limbs (Syphilinum)

II Bubo with pain in spot on middle of r. thigh in front, only when standing and on deep pressure, which seemed to touch spot, which was apparently on periosteum.

II For two or three winters intense cold pain in both legs, < in l., came on every night on lying down, lasting all night ; > by getting up and walking, and in warm weather.

II Osteosarcoma in centre of r. tibia the size of half an ostrich egg, pains agonizing at night, growth irregular, spongy, partly laminated, very hard.

II Redness and rawness with terrible itching between toes.

II Rheumatic swelling of left wrist and big toe, bluish red, with pains as if somebody sawed at his bones with a dull saw ; > by heat of stove ; < from sundown to sunrise ; no appetite ; has lasted two weeks.

II Severe attacks of aching in lower limbs.

II Sharp rheumatic pain, burning like fire, in 1, side of r. instep and below inner malleolus, prevents her from moving foot < when toe is pointed inward ; < in evening, continuing during' night, waking her up suddenly every two or three hours, worst from 1 :30 to 3 :30 A. M., < toward daybreak (improved).

II Slight contraction of tendons beneath r. knee.

II Two ulcers larger than a crown piece, dirty stinking, sloughing, with jagged, elevated edges, one on thigh above patella, another on head of tibia ; two large pieces of bone came away from head of tibia.

I Contracted, painful feeling in soles, as if tendons were too short.

I Excruciating arthritis ; swelling, heat, and redness intense.

I Rheumatism, muscles are caked in hard knots or lumps.

Skin (Syphilinum)

II A blood-boil on arm : face broken out with a lumpy fiery rash.

II Biting sensation in different parts of body, as if bitten by bugs, at night only.

II Eruption over whole body not elevated, but could be distinctly felt by passing hand over skin ; after Syphilinum im. eruption came rapidly to surface ; at same time a disagreeable odor began to be developed ; eruption reddish-brown like smallpox pustules, without central depression ; body covered with it, except

scrotum and penis ; increased, completely covering inside of mouth and throat, making it difficult to swallow even liquids ; eyes also covered, making him completely blind ; intolerable smell from body ; tips of pimples became filled with pus ; < from warmth of bed ; fetid breath ; eruption developed still more, a great quantity of pus, with intolerable itching, yet could not scratch as it was extremely sore ; eruption left skin of entire body covered with dull, reddish, copper-colored spots, which in cold, looked blue.

II Macula ; copper-colored ; from crown of head to sole of foot.

II Macula over back, chest, abdomen, arms and legs, but not on any uncovered part of body.

II Several elevated spots on arm, stomach, leg and finger ; has them habitually on face, chiefly on left cheek.

II Syphilitic bulla discharging freely on cheeks, under chin, on back of shoulders, on scalp and other parts of body (infantile syphilis).

II Syphilitic rash, very prominent on forehead, chin, arms and front of thorax, an abundance of fine scales peeling off ; large prominent spot on centre of forehead, filled with fluid, as also are some smaller patches.

I Pemphigus, looking like a pock, often confluent and persistently reappears.

Sleep (Syphilinum)

I Absolute sleeplessness (vies with Sul. in producing quiet, refreshing sleep).

Fever (Syphilinum)

II Fever from 11 to 1 P. M. daily ; perspires when she begins to get over fever ; pain in back, < between shoulders, no ambition or desire to move.

II Great pains in head, whole body extremely cold, looked blue ; wanted to be covered with blankets or couldn't get warm ; no appetite ; sleeping almost continually, could not be aroused.

II Nervous chills preceded by pains in head, esp. occiput and scalp of that part ; pains below waist, in pelvis, legs, esp. tibia, which is sensitive to touch ; bowels torpid ; cross, irritable, peevish : pains begin every day at 4 P. M., culminate at midnight, disappear at daylight.

II Sweat : profuse at night, sleepless and restless ; esp. between scapula and down to waist, with excessive general debility.

I Excessive general debility and continued night sweats, latter being most marked between scapula and down to waist .

Generalities (Syphilinum)

II Body, extremities and face covered with syphilides ; a sticking soreness begins in throat every evening between 6 and 7 o'clock and continues to grow < during night ; exceedingly restless until 4 A. M. ; then a restless sleep for a few hours ; can scarcely swallow ; when swallowing a sensation as of throat tearing to pieces ; continual throbbing in throat, < from cold and hot drinks, and < lying down ; throbbing in temples and ears, boring in ears meeting in centre of brain ; sensation as if top of head were coming off ; drawing pain in eyes < from lamplight ; teeth pain when eating, also when taking anything hot or cold, feel as if they were loose, pressing teeth together and pressing throat with hands ; excessive flow of saliva, it runs out of mouth when sleeping ; severe pain in neck ; bending back head, > pain in neck ; aching pain in shoulders and knees ; rending, tearing pains throughout body, > moving about slowly ; had his wife hide his revolver lest in a fit of desperation he might kill himself, as was his desire during extreme paroxysms of pain ; strikes wall with fist and beats head against wall for relief ; stools hard, dry like sheep dung ; desire for stool three or four times a day, but only a little scentless wind passes which gives relief ; sitting at stool, > pain in head ; is easily offended, gets desperate, cannot bear to be alone, great anxiety about getting well ; at night no position suits him, walks floor or goes into street and moves about slowly ; sleepy all the time but cannot

sleep ; dreams about his disease ; < in open air ; frequent urination with sudden desire : discharges large quantities of muddy urine.

o

TUBERCULINUM.

Mind (Tuberculinum)

IRRITABLE ON WAKING ; NOTHING CAN PLEASE HIM ; NOTHING SATISFIES. Nervous ; weak ; IRRITABLE.

II Although naturally of a sweet disposition, became taciturn, sulky, snappish, fretty, irritable, morose, depressed and melancholic, even to *insanity.-Burnett.*

II With every little ailment whines and complains ; easily frightened, particularly by dogs ; screams in terror when approached by a dog.

Head (Tuberculinum)

II Basilar meningitis. - Sinker.

II Boy, aged 20 months, ill for days with head, high fever, restlessness and constant screaming ; finally no sleep for forty hours, followed by a condition of collapse ; peculiar smell of body ; family history of tuberculosis.-**Burnett.**

II Fretful and ailing, whines and complains, indurated glands can be felt everywhere, child hot, drowsy, urine red and sandy, much given to be frightened, particularly by dogs ; was vaccinated and had a very bad arm for four months thereafter ; would not smile, whimpers when spoken to, skin dingy, skull hydrocephalic.-**Burnett.**

II Headache of forty-five years' standing, *pain passing from r. frontal protuberance to r. occipital region.-Swan.*

II Headache of great intensity preceded by a shuddering chill passing from brain down spine, with attack a feeling as if head above eyes were swollen ; became unconscious with screaming, tearing her hair, beating her head with her fists or trying to dash it against wall or floor.-**Swan.**

II Sullen, taciturn, irritable, screams in his sleep, is very restless at night, constipated ; sister died of tubercular meningitis. - **Burnett.**

II Terrible pain in head, as if he had a tight hoop of iron around it ; trembling of hands ; distressing sensation of damp clothes on his spine ; almost absolute sleeplessness ; profound adynamia ; was thought by his friends to be-on verge of insanity ; most of his brothers and sisters had died of water on brain ; r. lung solid, probably from healed-up cavities, as he at one time suffered from pulmonary *phthisis.-Burnett.*

II Tubercular meningitis, with effusion ; head gradually enlarged ; alternately wakeful and delirious at night, talked nonsense by day, at intervals ; nocturnal hallucinations and fright, delirium ; pyrexia ; had eczema which almost disappeared after two unsuccessful vaccinations, and which were soon followed by above condition ; after administration of remedy there occurred a severe pustular eruption, then patches of a lepra and eczema appeared. -**Burnett**

I Headache, *with frequent sharp, cutting pains passing from above r. eye through head to back of left ear.-Rose.*

Stomach (Tuberculinum)

II Windy dyspepsia, with pinching and pains under ribs of r. side in mammary line.-**Burnett.**

Abdomen (Tuberculinum)

II Fever, emaciation, abdominal pains and discomfort, restless at night, glands of both groins enlarged and indurated ; cries out in sleep ; strawberry tongue.-*Burnett*.

II Inguinal glands indurated and visible ; excessive sweats ; chronic diarrhea.

II Tabes mesenterica ; swelling on left side, also on r. ; complains of a stitch in side after running ; languid and indisposed to talk ; nervous and irritable ; talks in his sleep ; grinds his teeth ; appetite poor ; hands blue ; indurated and palpable glands everywhere ; a drum belly ; spleen region bulging out. -*Burnett*.

Stool And Anus (Tuberculinum)

II Cholera infantum. - *Swan*.

II Severe hemorrhage from bowels, cough ; emaciation ; family history of phthisis.

Mrs. S , aged 27, Denver, catarrh of nose, throat and./ larynx ; cough severe < by lying down ; for ten or fifteen minutes after retiring coughs severely, loosens some tough mucus which can neither be raised nor swallowed.

Night sweats if warmly covered. Morning diarrhea : watery, profuse, gushing, at 6 :30 A. M. ; a slight motion, turning in bed, necessitates getting up in a hurry, with a rush. Rumbling, gurgling in abdomen as if quarts of water were in stomach and bowels. By careful eating can go rest of day without. Going without noon-day meal stool is natural, but it would not affect morning diarrhea. In morning, weak ; exhausted. In after noon, feels fairly well. Excessively nervous during menses. Last three periods have been ten or fifteen days late and more. scanty than usual. Under the care of one of our best homeopathic prescribers has had Aloe, Sulphur, Podophyllum, Gamboge and Rumex.

Respiratory Organs (Tuberculinum)

II Expectoration of non-viscid, very easily detached, thick phlegm from air passages, followed after a day or two by a very clear ring of voice.-*Burnett*.

II Hard, dry cough, shaking patient, more during sleep, but did not waken him. -*Boardman*.

II Hard, dry cough, sometimes slight, but generally no expectoration, slightly feverish.-*Boardman*.

II Slight tedious hacking cough, which had lasted for months in a girl of a distinctly phthisic habit.-*Burnett*.

Chest (Tuberculinum)

II A nasty little cough, for seven weeks ; much expectoration : pains in r. lung ; evening fever ; liver and spleen enlarged ; cough morning after breakfast ; neck slightly goitrous ; eats hardly any *breakfast*.-*Burnett*.

II Anemic, sickly, pale ; profound debility ; dyspnea, cannot mount or hurry ; menses *irregular*.-*Burnett*.

II Bad cough of about twelve months' duration ; expectoration of blood ; one of apices was audibly diseased ; has had pneumonia ; chest flat ; respiration accelerated ; tanned unduly in *sun*.-*Burnett*.

II Complains that she has been in consumption for many years ; is very thin and consumed with fever ; lungs very flat ; respiration almost imperceptible ; fever ; poor appetite ; languid.-*Burnett*.

II Cough, < 6 A. m. ; notched incisors ; thin and puny ; cervical and inguinal glands much enlarged and indurated ; strawberry tongue ; girl, aged 7.-*Burnett*.

II Girl, aged 15, tall for her age ; tonsils enlarged ; chronic discharge from nose, < early morning on rising ; speech thick ; thorax of pigeon-breast type ; perspires much across nose ; very bad perspiration of chest, armpits, palms, nose, and feet ; feels very chilly ; spleen swollen ; distinct dullness on percussion at apex of r. lung ; suffered badly from vaccination ; gets chilblains. -*Burnett*.

II Hectic flush of cheeks ; shortness of breath ; slight hacking cough ; several strumous scars on neck ; dusky skin ; large, moist rales in both lungs ; increased vocal resonance of r. lung ; amphoric sounds in r. lung ; large, soft-feeling gland in left side of neck ; very pronounced endocardial bruit, best heard at apex beat. Iodoformum 3x in four grain doses for two months, with improvement, followed by Tuberculinum c. in very infrequent *doses.-Burnett.*

II Incipient phthisis in a boy aged 7 ; loss of flesh ; great prostration ; morbid timidity ; glands of groins and on both sides of neck very much enlarged and indurated, particularly glands over apex of r. lung ; as he had suffered much from vaccination, Thuja 30 and Sabina 30 were first given, then *Tuberculinum.-Burnett.*

II Incipient tubercular disease ; restless at nights ; sleepless ; grinds teeth ; tendency to diarrhea ; want of appetite ; foul breath ; notched teeth ; pain after food ; vomiting of food ; indurated glands ; strawberry tongue ; naughty ; very irritable temper ; puny growth ; very thin ; girl, aged 6.- *Burnett.*

II Lady, aged 26, in first stage of consumption ; dyspnea and rapid breathing ; loss of flesh ; greasy, dingy *skin.-Burnett.*

II Much fever, < evenings ; restless and terribly irritable ; much depressed and in almost constant agitation ; tongue very red ; chronic diarrhea ; has lost fourteen pounds during last six weeks ; has no appetite ; evacuations discharged from bowels as from *a pop-gun.-Burnett.*

II No respiratory sounds at top of r. lung, and vocal resonance slightly increased ; pain in left side ; profuse perspiration ; girl, aged 18.-*Burnett.*

II Nocturnal perspirations ; notched incisors ; indurated glands everywhere, very large and numerous ; drumbellied ; grinding of teeth at night ; great susceptibility to taking cold ; perspiration < at back of lungs and on head ; big head, with bulging forehead ; subject to attacks of fever and diarrhea. *Burnett.*

II Pronounced phthisical habit ; severe piles ; constipation ; brown cutaneous affection on *abdomen.-Burnett.*

II Ringworm on scalp ; lymphatic glands everywhere palpable ; ribs very flat ; strawberry tongue, bad cough, < at night ; although 11 years old she had practically no teeth, they were rudimentary, and not above level of *gums.-Burnett.*

II Slight hacking cough, continuing all day, < at bedtime and on rising ; emaciation ; dullness on percussion at apex of r. lung.-*Burnett.*

II Stout man, bright, florid complexion, mother died of phthisis, with which disease her sister is suffering ; gets pneumonia very often in cold weather ; hence travels from place to place to avoid colds ; coughs much, brings up much phlegm ; perspired profusely and drank great quantities of fluids ; wretched sleepless nights, with almost constant fever ; glands of neck much enlarged. *-Burnett.*

Neck And Back (Tuberculinum)

II Indurated cervical glands.

Lower Limbs (Tuberculinum)

II Tubercular swelling of knee ; intermittent attacks of pain in it ; has expectorated clots of blood and suffered from exhausting sweats ; family history of *phthisis.-Burnett.*

II Tuberculous disease of left knee ; for eleven months had been limping ; knee much enlarged and very tender ; teeth dirty and carious ; strawberry *tongue.-Burnett.*

Skin (Tuberculinum)

II Very bad tempered ; very much pigmented where sun's rays impinged upon him ; teeth dirty, *greenish.-Burnett.*

I Ringworm.

Fever (Tuberculinum)

II Fever.

o **USTILAGO.**

Mind (Ustilago)

II Very sad, cries frequently ; exceedingly prostrated from sexual abuse and loss of semen ; sleep restless.

Head (Ustilago)

I Nervous headache from menstrual irregularities in nervous women.

I Vertigo at climaxis with too frequent and profuse menstruation.

Male Sexual Organs (Ustilago)

II Spermatorrhea after onanism ; emissions every night, talking about women causes an emission ; very sad ; cries frequently ; says he cannot break off habit, has no control of himself when passion is aroused ; knows it is fast killing him ; cannot work, is so prostrated.

I Erotic fancies.

I Seminal emissions and irresistible tendency to masturbation.

Female Sexual Organs (Ustilago)

II Every day from 12 M. to about 4 P. M., constant pain from l. ovary to uterus ; every few minutes, pain in ovary is intensely severe, cutting like a knife ; pain in r. ovary and hypogastric region, but all starting from left ovary ; ovary can be distinctly felt in groin about as large as a hen's egg and very hard ; when pressed upon gives intense pain ; every day thinks she has fever with paroxysm of pain, but no chill ; slight leucorrhoea ; loss of appetite ; constipated.

II Extreme pain during period ; flow very profuse and did not cease entirely until next period ; most of time confined to bed.

II For days oozing of dark blood with small coagula ; uterus enlarged, cervix tumefied or dilated.

II Intermittent neuralgia of left ovary ; enlarged, very tender to touch.

II Lochia too profuse, partly fluid, partly clotted ; prolonged bearing-down pains ; uterus feels drawn into a knot.

II Menses every three weeks, with dark coagulum ; profuse, with gushes of bright-red blood when rising from a seat, or after having been startled or frightened ; two days before menses, a heavy backache with sharp pain across abdomen from hip to hip, followed by expulsive pains ; pains diminish after flow commences and stop with it ; between menstrual periods heavy dragging backache on exertion ; pain shooting up back from hips to shoulder ; abdomen tender to touch ; excessive bearing down ; pressure in head ; sensation of contraction in vertex, and feeling as if head were lifting off ; vertigo ; excoriating, albuminous leucorrhoea, < before menses ; ravenous appetite ; excessive tired feeling ; pulse 80 and weak ; mental depression.

II Menses suppressed for last fourteen months ; very irritable and depressed ; uneasiness in region of stomach ; pain in ovarian region, especially left ; skin hot and dry ; constipation, stools dry and hard ; no

appetite ; stitching pains in chest, < worse in left side ; constant hacking cough ; considerable expectoration ; night sweats ; general prostration ; great uneasiness in lower extremities.

II Mild leucorrhoea.

II Oozing of dark blood, highly coagulated, forming occasionally long, black, stringy clots.

II Ovarian irritation, constant pain in left ovary passing down hip, has to limp when walking ; pains sharp and at times pass down leg with great rapidity ; every few days has quite a swelling in left groin ; cannot bear pressure over ovary.

II Ovaritis ; took cold after menstruation ; constant dull pain in r. groin and back, three or four times an hour ; sharp neuralgic pains in ovary ; walking painful ; bowels torpid, very languid.

II Profuse discharge of dark, clotted blood of fetid odor, with pain and tenderness in one or both ovaries.

II Profuse menstruation, flow lasting from ten days to two weeks, at first very abundant, gradually wearing off ; always < from motion ; discharge dark and quite painless.

II Severe menorrhagia for past twelve years at every menstrual period, lasting a week or ten days, sometimes longer ; pale, thin, weak, very nervous.

II Sub serous or interstitial fibroid of uterus (two cases), fibroid much diminished.

II Subject to headaches ever since menstruation appeared at age of fifteen ; headache mostly on top of head ; appetite poor ; pain in left chest with some cough ; total suppression of menses for last eight months ; severe pain in back, is unable to ride in carriage ; pain in uterine region, especially over ovarian region, < left side ; vomiting of mucus and blood daily ; no sleep ; some leucorrhoea ; hysterical ; no uterine displacement, but great congestion in pelvic region.

II Subject to profuse menstruation ; childless ; large, fleshy, flabby, bloated-looking, with a very sallow complexion, inclined to be (and formerly had been) dropsical from excessive loss of blood ; profuse menstruation, which seems to her to be principally water and clots ; says there is no outward flow when she lies still, but clots and water pass out of uterus when she gets up ; feels so full in uterus that she must rise to get rid of clots ; flowed fearfully during night ; very low, scarcely able to speak aloud.

II Uterus hypertrophied, sensitive, blood bright, fresh, without coagula.

II Vicarious menstruation from lungs and bowels.

I Blood dark, but so thin as to scarcely color fingers.

I Cervix tumefied, bleeds when touched.

I Chronic uterine hemorrhage, and passive congestion.

I Climaxis : vertigo ; frequent flushing ; metrorrhagia.

I Constant aching distress at mouth of womb.

I Deficient labor pains ; os soft, pliable, dilatable.

I Dysmenorrhoea of a congestive character, with much ovarian irritation ; severe pain in ovaries ; uterus and back every few minutes ; scanty, pale flow accompanied by false membranes ; poor appetite, thickly-coated tongue.

I Hypertrophy and subinvolution of uterus with great atony.

I Menorrhagia at climaxis ; active and constant flowing with frequent clots.

I Menorrhagia, with displaced uterus.

I Metrorrhagia after miscarriage, confinement or at the climaxis.

I Suppression of menses without apparent cause ; troublesome cough ; considerable expectoration ; sometimes also dry cough ; stitching pains in chest, especially left side ; night sweats ; loss of appetite ; pain in ovaries, especially 1 ; general debility, headache ; leucorrhoea ; chlorotic ; anemic, as if in first stage of consumption.

I Uterus : hypertrophied ; prolapsed ; cervix sensitive, spongy.

Pregnancy. Parturition. Lactation (Ustilago)

II Agalactia ; chronic inflammation, and induration of mamma.

II Discharge of blood from uterus, bright-red, partly fluid, partly clotted ; passive congestion of uterus, so that there is a slight oozing of blood after each examination ; tissues of uterus feel soft and spongy ; os patulous.

II For last year vertigo every day, some days so bad she has to go to bed ; menses every three weeks for last year ; last about ten days and profuse ; flows as much again as she did when in health ; constant aching distress under left mamma ; rheumatic pains in shoulders and back ; very weak, not able to work.

II Has aborted a number of times at third month ; is now about three months pregnant ; for last ten days has had more or less hemorrhage every day, some days quite bad ; not so much at night ; blood passes a number of times through day, in dark-colored clots.

II One and a half hours after delivery commenced to flow violently.

II Promotes expulsion of foreign bodies from the uterus.

II Puerperal peritonitis ; aborted about two days since, at about three months ; constant fever ; pulse 120 ; cannot bear least pressure on any portion of bowels ; about six times today has had sharp, cutting pains in left ovary ; has flowed constantly for two days ; blood dark, not copious, nor attended with bearing-down pains ; cannot move in bed ; is compelled to lie upon her back ; constant, dull, frontal headache ; loss of appetite, tongue furred.

II Severe flooding two weeks after labor ; large bright-red clots ; no pain ; very weak.

Very profuse lochial discharge, very dark in color, almost black.

I Constant flooding ; every few minutes, expulsion of a large clot of bright red blood, with bearing-down pains.

I Fibroids and induration of os.

I Passive hemorrhage after miscarriage, blood in lumps, flooding for days and weeks.

I Persistent hemorrhage of brownish blood, with want of uterine contraction.

I Post-partum hemorrhages from a flabby, atonic condition of uterus.

I Puerperal peritonitis, with constant flooding ; high fever ; secretion putrid ; abdomen excessively tender and tympanitic.

Upper Limbs (Ustilago)

I Hypertrophy or loss of nails.

o **VARIOLINUM.**

Eyes (Variolinum)

II Chronic ophthalmia with loss of sight.

II Pupils contracted.

Mouth (Variolinum)

II Thick, dirty yellow coating on tongue.

II When asleep tongue protruded, black coating, when raised it is with difficulty drawn back ; looks like a mass of putrid flesh.

I Tongue coated white, as with a piece of white velvet, in variola with headache, backache and high fever.

Throat (Variolinum)

II Pharynx and fauces deep purplish crimson, with gangrenous appearance ; breath horribly offensive.

Stomach (Variolinum)

II Frequent bilious vomiting.

Stool (Variolinum)

II Dysentery.

Respiratory Organs (Variolinum)

II Asthma.

Lower Limbs (Variolinum)

II Severe chill followed by high fever ; severe pain in back as if broken ; pain all over head, very severe and constant in occiput ; frequent bilious vomiting ; thick, dirty, yellowish coating on tongue ; wild delirium and spasms ; night before eruption appeared obstinate constipation ; on third day very thick eruption of small-pox pustules, soon assuming confluent form, Variolin. cmm.

II Small-pox ; peculiar smell causes intense sickness of stomach ; congestion toward head, palpitation of heart ; two hours after took a dose Variolin. 30 ; 'an hour later crawling in back and feeling of coldness in lower extremities ; cold feet ; lame, heavy feeling in left arm ; no appetite ; sleep disturbed by heat ; toward morning some perspiration over body ; next day continuous pressing headache, especially in occiput ; pulse somewhat irritated ; disinclined to mental work ; when reading heat in head and forehead ; weak when walking ; pain in joints of upper and lower extremities, as if lame ; toward evening again feverish ; pressing pain in small of back down to sacrum (Glonoin. 3 relieved headache) ; in bed, drawing in upper jaw and teeth ; sleep full of dreams ; toward morning again perspiration ; urine smells ammoniacal ; on fourth day red pimples on back of left hand, staying for several days without filling with pus ; mental work still causes heat in forehead and pressure in head ; escaped smallpox.

II Terrible pains in back on r. side of spine, and over and below shoulder blade ; muscles sore to touch, nausea, pains all over especially in legs ; tongue clean ; pulse 120 ; Variolin. cm. ; body completely covered with large pustules, face one mass of confluent pustules, pulse still high, constant expectoration of viscid

mucus, mouth and fauces lined with pustules, even tongue covered with them ; bowels constipated, mild delirium at times ; eight days later temperature 104/ ; pulse 120, very weak and stopping at intervals ; great fear of death ; begging to know if he must die, and before sentence was completed would drop into a heavy sleep with stertorous breathing, jaw dropped on breast, pupils contracted, teeth covered with thick brown slime, centre of tongue perfectly black, mucous membrane of mouth and pharynx of a deep purplish-crimson, with gangrenous appearance and breath horribly offensive ; skin of face and neck of a deep dark purple ; odor from body like a fetid stream ; little control over tongue or jaw, latter hanging down, and tongue protruding like a mass of decayed liver when asleep ; an effort to speak when roused up caused violent trembling of jaw and tongue, which was drawn back into mouth with difficulty, was stiff, but looked like a mass of putrid flesh ; urine dark colored, passing freely through whole attack ; had continued the cm. till now, gave one dose cmm. dry on tongue ; next day almost convalescent ; made a good recovery with but few marks.

I On third day of eruption, when pustules had filled, and were confluent on face, intense itching ; Variolinum 1m. in water, every two hours ; on second day itching nearly gone, pustules shrinking ; third day drying ; fifth day crusts fell from face ; seventh day other crusts fell off, and tongue clean.

Have used and seen it used in many and severe cases, and when treated with Variolin. 200 disease is shortened nearly or quite one-half, sufferings of patients much mitigated, secondary fever either absent or very much lighter, pustules do not burst, but wither or wilt and fall off, suppurative stage immeasurably hastened and shortened, and patients are not marked.
